## KDM4D Antibody

Catalog No: #36563



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


	Accr	INTIAN	
u	COUL	iption	

Product Name	KDM4D Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antigen affinity purification.
Applications	WB IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous levels of total KDM4D protein.
Immunogen Type	Recombinant Protein
Immunogen Description	Fusion protein corresponding to a region derived from internal residues of human lysine (K)-specific
	demethylase 4D
Target Name	KDM4D
Other Names	JMJD2D
Accession No.	Swiss-Prot#: Q6B0I6NCBI Gene ID: 55693Gene Accssion: BC122858
Uniprot	Q6B0I6
GeneID	55693;
SDS-PAGE MW	59kd
Concentration	2.2mg/ml
Formulation	Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
Storage	Store at -20°C

## Application Details

Western blotting: 1:500-1:2000
Immunohistochemistry: 1:50-1:200

## **Images**




Gel: 8%SDS-PAGE

Lysates (from left to right): Human fetal brain tissue

Amount of lysate: 40ug per lane Primary antibody: 1/1100 dilution Secondary antibody dilution: 1/8000

Exposure time: 10 minutes



Immunohistochemical analysis of paraffin-embedded Human lung cancer tissue using #36563 at dilution 1/50.

## Background

JMJD2D (Jumonji domain-containing protein 2D), also known as JHDM3D or KDM4D, is a 520 amino acid protein that belongs to the JHDM3 histone demethylase family. Localized to the nucleus, JMJD2D functions as a histone demethylase that removes specific methyl residues from Histone H3, thereby playing a crucial role in the histone code. JMJD2D binds iron as a cofactor and contains one JMJC domain and one JMJN domain, both of which are thought to exhibit enzymatic activity during chromatin remodeling events. In addition, JMJD2D forms a complex with the ligand-bound form of the androgen receptor (AR) and, through this interaction, activates AR expression. Overexpression of AR is associated with prostate cancer, suggesting that, via its ability to upregulate AR, JMJD2D may be involved in carcinogenesis.

Note: This product is for in vitro research use only