

Tau(Phospho-Thr205) Antibody

Catalog No: #11108

Package Size: #11108-1 50ul #11108-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	Tau(Phospho-Thr205) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of Tau only when phosphorylated at threonine 205.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 205 (P-G-T(p)-P-G) derived from Human Tau.
Conjugates	Unconjugated
Target Name	Tau
Modification	Phospho
Other Names	MAPT; MTAPT; MTBT1; Neurofibrillary tangle protein; PHF-tau
Accession No.	Swiss-Prot: P10636NCBI Protein: NP_001116538.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 48 62 78 kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from mouse brain tissue using Tau(Phospho-Thr205) Antibody #11108 and the same antibody preincubated with blocking peptide.



Immunohistochemical analysis of paraffin-embedded rat hippocampal region tissue from a model with Alzheimer

Background

Promotes microtubule assembly and stability, and might be involved in the establishment and maintenance of neuronal polarity. The C-terminus binds axonal microtubules while the N-terminus binds neural plasma membrane components, suggesting that tau functions as a linker protein between both. Axonal polarity is predetermined by tau localization (in the neuronal cell) in the domain of the cell body defined by the centrosome. The short isoforms allow plasticity of the cytoskeleton whereas the longer isoforms may preferentially play a role in its stabilization.

del C Alonso A, et al. (2004) J Biol Chem; 279(33): 34873-81

Li G, Yin H, Kuret J (2004) J Biol Chem; 279(16): 15938-45

Giasson BI, et al. (2002) Biochemistry; 41(51): 15376-87

Liu F, et al. (2002) FEBS Lett; 530(1-3): 209

Taniguchi T, et al. (2001) J Biol Chem; 276(13): 10025-31

Published Papers

Tan Wenfei, Cao Xuezhao, Wang Junke et al., Tau hyperphosphorylation is associated with memory impairment after exposure to 1.5% isoflurane without temperature maintenance in rats., European Journal of Anaesthesiology, 27(9):835-841. doi: 10.1097/EJA.0b013e32833a6561(2010)

[PMID:20485178](#)

el at., Trillium tschonoskii maxim extract attenuates abnormal Tau phosphorylation.Neural Regen Res.In Neural Regen Res.On 2018 May by Luo HB, Shang N. et al..PMID: 29863023, , (2018)

[PMID:29863023](#)

el at., Liraglutide Ameliorates Hyperhomocysteinemia-Induced Alzheimer-Like Pathology and Memory Deficits in Rats via Multi-molecular Targeting. In Neurosci Bull on 2019 Jan 10 by Zhang Y, Xie JZ, et al..PMID: 30632006, , (2019)

[PMID:30632006](#)

el at., Tau-Induced Ca²⁺/Calmodulin-Dependent Protein Kinase-IV Activation Aggravates Nuclear Tau Hyperphosphorylation .In Neurosci Bull. On 2018 Apr by Wei YP, Ye JW et al..PMID: 28646348, , (2018)

[PMID: 28646348](#)

el at., Adiponectin Attenuates Streptozotocin-Induced Tau Hyperphosphorylation and Cognitive Deficits by Rescuing PI3K/Akt/GSK-3 β Pathway. In Neurochem Res.On 2018 Feb by Xu ZP, Gan GS et al..PMID: 29098530, , (2018)

[PMID: 29098530](#)

el at., Tau hyperphosphorylation is associated with memory impairment after exposure to 1.5% isoflurane without temperature maintenance in rats.In Eur J Anaesthesiol on 2010 Sep by Tan W, Cao X, et al..PMID:20485178, , (2010)

[PMID:20485178](#)

el at., Protective effects of lithium treatment for spatial memory deficits induced by tau hyperphosphorylation in splenectomized rats.In Clin Exp Pharmacol Physiol on 2010 Oct by Tan WF, Cao XZ,et al..PMID:20659131, , (2010)

[PMID:20659131](#)

el at., Neuroglobin attenuates Alzheimer-like tau hyperphosphorylation by activating Akt signaling.In J Neurochem on 2012 Jan by Chen LM, Xiong YS,et al..PMID:21496024, , (2012)

[PMID:21496024](#)

el at., Synaptic released zinc promotes tau hyperphosphorylation by inhibition of protein phosphatase 2A (PP2A). In J Biol Chem on 2012 Mar 30 by Xu-Ying Sun, Yu-Ping Wei, et al..PMID: 22334661, , (2012)

[PMID:22334661](#)

el at., Cnga2 Knockout Mice Display Alzheimer's-Like Behavior Abnormities and Pathological Changes .In Mol Neurobiol on 2016 Sep by Ao-Ji Xie, En-Jie Liu et al..PMID: 26377105, , (2016)

[PMID:26377105](#)

el at., Olfactory Deprivation Hastens Alzheimer-Like Pathologies in a Human Tau-Overexpressed Mouse Model via Activation of cdk5.In Mol Neurobiol.On 2016 Jan by Li K, Liu FF et al..PMID:25465240, , (2016)

[PMID:25465240](#)

el at., Brain-derived neurotrophic factor protects against tau-related neurodegeneration of Alzheimer's disease.In Transl Psychiatry on 2016 Oct 4 by S-S Jiao , L-L Shen et al..PMID:27701410 , , (2016)

[PMID:27701410](#)

el at., Region-specific expression of tau, amyloid- ϵ Y protein precursor, and synaptic proteins at physiological condition or under endoplasmic reticulum stress in rats.In J Alzheimers Dis on 2014 by Li Lin, Shu-Sheng Yang, et al..PMID: 24787918, , (2014)

[PMID:24787918](#)

el at., Extrasynaptic NMDA Receptor-Induced Tau Overexpression Mediates Neuronal Death Through Suppressing Survival Signaling ERK Phosphorylation.In Cell Death Dis on 2016 Nov 3 by Xu-Ying Sun , Qing-Zhang Tuo et al..PMID: 27809304, , (2016)

[PMID:27809304](#)

el at., Interneuron Accumulation of Phosphorylated tau Impairs Adult Hippocampal Neurogenesis by Suppressing GABAergic Transmission. In Cell Stem Cell on 2020 Mar 5 by Jie Zheng, Hong-Lian Li,et al..PMID:31978364, , (2020)

[PMID:31978364](#)

el at., Genetic deletion of $\alpha 7$ nicotinic acetylcholine receptors induces an age-dependent Alzheimer's disease-like pathology. In Prog Neurobiol on 2021 Nov by Maria Rosaria Tropea, Domenica D Li Puma,et al..PMID:34453977, , (2021)

[PMID:34453977](#)

el at., A novel dephosphorylation targeting chimera selectively promoting tau removal in tauopathies. In Signal Transduct Target Ther on 2021 Jul 14 by Jie Zheng, Na Tian et al..PMID:34262014, , (2021)

[PMID:34262014](#)

el at., Effectiveness and mechanisms of combined use of antioxidant nutrients in protecting against oxidative stress-induced neuronal loss and related neurological deficits. In CNS Neurosci Ther on 2024 Jul by Lu Wang, Yingjuan Wei,et al..PMID:39072940, , (2024)

[PMID:39072940](#)

Yanmin Chang;Yi Yao;Rong Ma;Zemin Wang;Junjie Hu;Yanqing Wu;Xingjun Jiang;Lulu Li;Gang Li el at., DI-3-n-Butylphthalide Reduces Cognitive Deficits and Alleviates Neuropathology in P301S Tau Transgenic Mice, , (2021)

[PMID:](#)

Xing Jun Jiang;Yan Qing Wu;Rong Ma;Yan Min Chang;Lu Lu Li;Jia Hui Zhu;Gong Ping Liu;Gang Li el at., PINK1 Alleviates Cognitive Impairments via Attenuating Pathological Tau Aggregation in a Mouse Model of Tauopathy, , (2022)

[PMID:](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.