## Zap-70(Phospho-Tyr319) Antibody

Catalog No: #11159

Description

Package Size: #11159-1 50ul #11159-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Product Name          | Zap-70(Phospho-Tyr319) Antibody                                                                        |
|-----------------------|--------------------------------------------------------------------------------------------------------|
| Host Species          | Rabbit                                                                                                 |
| Clonality             | Polyclonal                                                                                             |
| Purification          | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.       |
|                       | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho |
|                       | specific antibodies were removed by chromatogramphy using non-phosphopeptide.                          |
| Applications          | WB IHC IF                                                                                              |
| Species Reactivity    | Hu Ms Rt                                                                                               |
| Specificity           | The antibody detects endogenous level of VEGFR2 only when phosphorylated at tyrosine 1214.             |
| Immunogen Type        | Peptide-KLH                                                                                            |
| Immunogen Description | Peptide sequence around phosphorylation site of tyrosine 319 (S-P-Y(p)-S-D) derived from Human Zap-70. |
| Conjugates            | Unconjugated                                                                                           |
| Target Name           | Zap-70                                                                                                 |
| Modification          | Phospho                                                                                                |

## **Application Details**

Other Names

Accession No.
Calculated MW

Concentration

Formulation

Storage

Western blotting: 1:500~1:2000 Immunohistochemistry: 1:50~1:200 Immunofluorescence: 1:100~1:200

SRK, STD, TZK, STCD, ZAP-70


sodium azide and 50% glycerol.

Predicted MW: 70kd

1.0mg/ml

Swiss-Prot: P43403NCBI Protein: NP\_001070.2


## **Images**




Western blot analysis of extracts from Jurkat cells using Zap-70(Phospho-Tyr319) Antibody #11159(Lane 2) and the same antibody preincubated with blocking peptide(Lane1).

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%

Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.



Immunofluorescence staining of methanol-fixed Hela cells using Zap-70(Phospho-Tyr319) Antibody #11159.



Western blot analysis of extracts from HL60 cells, treated with calf intestinal phosphatase (CIP), using Zap-70 (Phospho-Tyr319) Antibody #11159.

## Background

Plays a role in T-cell development and lymphocyte activation. Essential for TCR-mediated IL-2 production. Isoform 1 induces TCR-mediated signal transduction, isoform 2 does not.

Salomon AR, et al. (2003) Proc Natl Acad Sci U S A; 100(2): 443-448

Ku GM, et al. (2001) EMBO J; 20(3): 457-465

Tang J, et al. (1999) Proc Natl Acad Sci U S A; 96(17): 9775-9780

Zhao Q, et al. (1996) Mol Cell Biol; 16(12): 6765-6774 Williams BL, et al. (1999) EMBO J; 18(7): 1832-1844

Adjali O, et al. (2005) J Clin Invest; 115(8): 2287-2295.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.