PLCg2(Phospho-Tyr753) Antibody

Catalog No: #11175

Package Size: #11175-1 50ul #11175-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

PLCg2(Phospho-Tyr753) Antibody
Rabbit
Polyclonal
Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
specific antibodies were removed by chromatogramphy using non-phosphopeptide.
WB IF
Hu Ms Rt
The antibody detects endogenous level of PLCg2 only when phosphorylated at tyrosine 753.
Peptide-KLH
Peptide sequence around phosphorylation site of tyrosine 753 (S-L-Y(p)-D-V) derived from Human PLCg2.
Unconjugated
PLCg2
Phospho
PLC-IV; PLC-gamma2; Phospholipase C-gamma-2
Swiss-Prot: P16885NCBI Protein: NP_002652.2
1.0mg/ml
Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
sodium azide and 50% glycerol.
Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 150kd


Western blotting: 1:500~1:1000

Immunofluorescence: 1:100~1:200

Images



Western blot analysis of extracts from HepG2 cells untreated or treated with EGF using PLCg2(Phospho-Tyr753) Antibody #11175.



Immunofluorescence staining of methanol-fixed Hela cells using PLCg2(Phospho-Tyr753) Antibody #11175.

Background

The production of the second messenger molecules diacylglycerol. (DAG) and inositol 1,4,5-trisphosphate (IP3) is mediated by activated phosphatidylinositol-specific phospholipase C enzymes. It is a crucial enzyme in transmembrane signaling.

Kim YJ, et al. (2004) Mol Cell Biol 24: 9986-9999

Humphries LA, et al. (2004) J Biol Chem 279 : 37651-37661 Suzuki-Inoue K, et al. (2004) Biochem J 378 : 1023-1029 Rodriguez R, et al. (2003) Biochem J 374 : 269-280

Note: This product is for in vitro research use only and is not intended for use in humans or animals.