

MEK1/MEK2(Phospho-Ser217/Ser221) Antibody

Catalog No: #11205

Package Size: #11205-1 50ul #11205-2 100ul

Description

Product Name	MEK1/MEK2(Phospho-Ser217/Ser221) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	IHC WB
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of MEK1/MEK2 only when phosphorylated at serine 217/221.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 217/221 (I-D-S(p)-M-A) derived from Human MEK1/MEK2.
Conjugates	Unconjugated
Target Name	MEK1/MEK2
Modification	Phospho
Other Names	ERK activator kinase 1/ERK activator kinase2; MAP kinase kinase 1/MAP kinase kinase 2; MAP2K1/MAP2K1; MAPK/ERK kinase 1/MAPK/ERK kinase 2; MAPKK 1/MAPKK 1
Accession No.	Swiss-Prot: Q02750NCBI Protein: NP_002746.1 NP_109587.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 45kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from HeLa cell untreated or treated with UV MEK1/MEK2(Phospho-Ser217/Ser221) Antibody#11205



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using MEK1/MEK2(Phospho-Ser217/Ser221) Antibody #11205(left) or the same antibody preincubated with blocking peptide(right).

Background

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates ERK1 and ERK2 MAP kinases.

Shen R, et al. (2002) *Mol Cell Biol*; 22(10): 3230-3236

Preisinger C, et al. (2005) *EMBO J*; 24(4): 753-765

Laine P, et al. (2000) *Biochem J*; 349(Pt 1): 19-25

Yaglom J, et al. (2003) *Mol Cell Biol*; 23(11): 3813-3824

Dahan S, et al. (2002) *Infect Immun*; 70(5): 2304-2310

Published Papers

el at., G protein subunit δ α regulates gastric cancer growth via the p53/p21 and MEK/ERK pathways. *In Oncol Rep* on 2017 Apr by Yizhuo Wang , Huijie Xiao, et al.. PMID: 28350126 , (2017)

[PMID:28350126](#)

el at., Response of MAPK pathway to iron oxide nanoparticles in vitro treatment promotes osteogenic differentiation of hBMSCs. *In Biomaterials*. On 2016 Apr by Wang Q, Chen B et al.. PMID:26874888 , (2016)

[PMID:26874888](#)

el at., Integrin δ α 6/Akt/Erk signaling is essential for human breast cancer resistance to radiotherapy. *In Sci Rep* on 2016 Sep 14 by Ting Hu, Rui Zhou et al.. PMID:27624978 , (2016)

[PMID:27624978](#)

el at., Effect of Chaihushugan San on Expression of the Raf/mitogen-activated Protein Kinase/Extracellular Signal-Regulated Kinase Pathway in the Hippocampi of Perimenopausal Rats Induced by Immobilization Stress . *In J Tradit Chin Med* on 2015 Aug by Li Shengqiang, Liang Wenna et al.. PMID:26427116 , (2015)

[PMID:26427116](#)

el at., MicroRNA δ α 46b, a sensitive indicator of mesenchymal stem cell repair of acute renal injury. *In Stem Cells Transl Med* on 2016 Oct by Yuan Zhu, Jing Yu et al.. PMID:27400799 , (2016)

[PMID:27400799](#)

el at., Suppressive action of acetate on interleukin-1 β production via tubulin- α 2-acetylation. In Immunol Cell Biol on 2014 Aug by Kazuhiro Ishiguro, Takafumi Ando et al..PMID:24777307, , (2014)

[PMID:24777307](#)

el at., Exosomes Derived From Human Mesenchymal Stem Cells Confer Drug Resistance in Gastric Cancer. In Cell Cycle on 2015 Aug 3 by Runbi Ji, Bin Zhang et al..PMID:26091251, , (2015)

[PMID:26091251](#)

el at., Mycoplasma ovipneumoniae induces sheep airway epithelial cell apoptosis through an ERK signalling-mediated mitochondria pathway. In BMC Microbiol on 2016 Sep 23 by Yanan Li, Zhongjia Jiang,et al..PMID:27663303, , (2016)

[PMID:27663303](#)

el at., TMEM16A-inhibitor loaded pH-responsive nanoparticles: A novel dual-targeting antitumor therapy for lung adenocarcinoma. In Biochem Pharmacol

on 2020 Aug by Shuai Guo, Liang Qiu,et al..PMID:32492446, , (2020)

[PMID:32492446](#)

el at., Inhibition of TMEM16A by Natural Product Silibinin: Potential Lead Compounds for Treatment of Lung Adenocarcinoma. In Front Pharmacol on 2021 Apr 14 by Shuai Guo, Xue Bai, et al..PMID:33935737, , (2021)

[PMID:33935737](#)

el at., TMEM16A, a Homoharringtonine Receptor, as a Potential Endogenic Target for Lung Cancer Treatment. In Int J Mol Sci on 2021 Oct 10 by Shuai Guo, Xue Bai, et al..PMID:34681590, , (2021)

[PMID:34681590](#)

el at., Matairesinoside, a novel inhibitor of TMEM16A ion channel, loaded with functional hydrogel for lung cancer treatment. In Int J Biol Macromol on 2024 Apr by Zhichen Wang, Ruili Geng, et al..PMID:38447844, , (2024)

[PMID:38447844](#)

el at., 7 β -dehydrocholesterol suppresses melanoma cell proliferation and invasion via Akt1/NF κ B signaling, In Oncol Lett on 2020 Dec by Jia Liu, Feiliang Zhong, et al..PMID: 33193858, , (2020)

[PMID:33193858](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.