Product Datasheet

Pyk2(Phospho-Tyr402) Antibody

Catalog No: #11216

Package Size: #11216-1 50ul #11216-2 100ul



Support: tech@signalwayantibody.com

Description **Host Species**

Pyk2(Phospho-Tyr402) Antibody **Product Name**

Rabbit

Clonality Polyclonal

Purification Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.

Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho

specific antibodies were removed by chromatogramphy using non-phosphopeptide.

WB IHC IF Applications Species Reactivity Hu Ms Rt

Specificity The antibody detects endogenous level of Pyk2 only when phosphorylated at tyrosine 402.

Peptide-KLH Immunogen Type

Peptide sequence around phosphorylation site of tyrosine 402 (D-I-Y(p)-A-E) derived from Human Pyk2. Immunogen Description

Unconjugated Conjugates

Target Name Pyk2

Modification Phospho

Other Names FADK 2; FAK2; Focal adhesion kinase 2; PTK2B; Proline-rich tyrosine kinase 2 RAFTK

Swiss-Prot: Q14289NCBI Protein: NP_004094.3 Accession No.

Concentration 1.0mg/ml

Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% Formulation

sodium azide and 50% glycerol.

Storage Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 116kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from 293 cells untreated or treated with Serum using Pyk2(Phospho-Tyr402) Antibody #11216



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Pyk2(Phospho-Tyr402) Antibody #11216(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using Pyk2(Phospho-Tyr402) Antibody #11216.

Background

Involved in calcium induced regulation of ion channel and activation of the map kinase signaling pathway. May represent an important signaling intermediate between neuropeptide activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. Interacts with the SH2 domain of Grb2. May phosphorylate the voltage-gated potassium channel protein Kv1.2. Its activation is highly correlated with the stimulation of c-Jun N-terminal kinase activity. Involved in osmotic stress-dependent SNCA 'Tyr-125' phosphorylation.

Gluck SL, et al. (2004) J Clin Invest; 114(12): 1696-1699

Benzing T, et al. (2001) Proc Natl Acad Sci U S A; 98(17): 9784-9789

Tian D, et al. (2002) Mol Cell Biol; 22(8): 2650-2662 Lu Z, et al. (2001) Mol Cell Biol; 21(12): 4016-4031 Krishnan HH, et al. (2006) J Virol; 80(3): 1167-1180

Published Papers

Mineko Tomomura, Hiroya Hasegawa, Naoto Suda el at., Serum Calcium-decreasing Factor, Caldecrin, Inhibits Receptor Activator of NF- B Ligand (RANKL)-mediated Ca2 Signaling and Actin Ring Formation in Mature Osteoclasts via Suppression of Src Signaling Pathway., THE JOURNAL OF BIOLOGICAL CHEMISTRY, 287(22):17963-17974.(2012)

PMID:22461633

el at., Serum Calcium-decreasing Factor, Caldecrin, Inhibits Receptor Activator of NF-I-• B Ligand (RANKL)-mediated Ca2+ Signaling and Actin Ring Formation in Mature Osteoclasts via Suppression of Src Signaling Pathway. In J Biol Chem on 2012 May 25 by Mineko Tomomura, Hiroya Hasegawa, et al..PMID:22461633, , (2012)

PMID:22461633

el at., Potential Involvement of M1 Macrophage and VLA4/VCAM-1 Pathway in the Valvular Damage Due to Rheumatic Heart Disease. In Front Biosci (Landmark Ed) on 2024 Jun 18 by Shenglin Xian, Yuan Li,et al..PMID:38940032, , (2024)

PMID:38940032

Yuwen Sheng;Chong Qiao;Zhonghui Zhang;Xiaoke Shi;Linhan Yang;Ruiying Xi;Jialing Yu;Wanli Liu;Guolin Zhang;Fei Wang el at., Calcium Channel Blocker Lacidipine Promotes Antitumor Immunity by Reprogramming Tryptophan Metabolism., , (2025)

PMID:39585774

Note: This product is for in vitro research use only and is not intended for use in humans or animals.