

mTOR(Phospho-Ser2448) Antibody

Catalog No: #11221

Package Size: #11221-1 50ul #11221-2 100ul

Description

Product Name	mTOR(Phospho-Ser2448) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB IHC IF
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of mTOR only when phosphorylated at serine 2448.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of serine 2448 (T-D-S(p)-Y-S) derived from Human mTOR.
Conjugates	Unconjugated
Target Name	mTOR
Modification	Phospho
Other Names	FRAP; FRAP1; FRAP2; RAFT1;
Accession No.	Swiss-Prot: P42345NCBI Protein: NP_004949.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 289kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using mTOR(Phospho-Ser2448) Antibody #11221(left) or the same antibody preincubated with blocking peptide(right).




Immunofluorescence staining of methanol-fixed MCF cells using mTOR(Phospho-Ser2448) Antibody #11221.



Western blot analysis of extracts from 293 cells untreated or treated with PMA using mTOR(Phospho-Ser2448) Antibody #11221.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using mTOR (Phospho-Ser2448) Antibody #11221



Western blot analysis of extracts from HeLa cells, treated with IFNa or calf intestinal phosphatase (CIP), using mTOR(Phospho-Ser2448) Antibody #11221.

Background

Kinase subunit of both mTORC1 and mTORC2, which regulate cell growth and survival in response to nutrient and hormonal signals. mTORC1 is activated in response to growth factors or amino-acids. Amino-acid-signaling to mTORC1 is mediated by Rag GTPases, which cause amino-acid-induced relocalization of mTOR within the endomembrane system. Growth factor-stimulated mTORC1 activation involves AKT1-mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potentially activates the protein kinase activity of mTORC1. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-421', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. mTORC2 is also activated by growth factors, but seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'.

Albanell J, et al. (2007) Clin Transl Oncol. 9(8):484-93.

Huang JJ, et al.

Published Papers

Macias M, Blazejczyk M, Kazmierska P et al., Spatiotemporal Characterization of mTOR Kinase Activity Following Kainic Acid Induced Status Epilepticus and Analysis of Rat Brain Response to Chronic Rapamycin Treatment. , PLoS ONE, 8(5): e64455.

doi:10.1371/journal.pone.0064455(2013)

[PMID:23724051](#)

el at., 5-HT 2 receptor mediates high-fat diet-induced hepatic steatosis and very low density lipoprotein overproduction in rats. In Obes Res Clin Pract. On 2018 Jan - Feb by Li X, Guo K et al.. PMID: 27133527, , (2018)

[PMID:27133527](#)

el at., Epithelial Cells Attenuate Toll-Like Receptor-Mediated Inflammatory Responses in Monocyte-Derived Macrophage-Like Cells to Mycobacterium tuberculosis by Modulating the PI3K/Akt/mTOR Signaling Pathway. In Mediators Inflamm. On 2018 Sep by Yang Y, Sun Y et al.. PMID:30356420, , (2018)

[PMID:30356420](#)

el at., Immunohistochemical assessment of growth factor signaling molecules: MAPK, Akt, and STAT3 pathways In oral epithelial precursor lesions and squamous cell carcinoma. In Odontology on 2019 May 6 by Tashiro K, Oikawa M, et al.. PMID:31062130, , (2019)

[PMID:31062130](#)

el at., Immunohistochemical assessment of growth factor signaling molecules: MAPK, Akt, and STAT3 pathways in oral epithelial precursor lesions and squamous cell carcinoma. In Odontology on 2020 Jan; by Tashiro K, Oikawa M, et al.. PMID: 31062130, , (2020)

[PMID:31062130](#)

el at., 5-Aza-2'-deoxycytidine inhibited PDGF-induced rat airway smooth muscle cell phenotypic switching. In Arch Toxicol on 2013 May by Ning Y, Huang H, et al.. PMID: 23423710, , (2013)

[PMID:23423710](#)

el at., Spatiotemporal Characterization of mTOR Kinase Activity Following Kainic Acid Induced Status Epilepticus and Analysis of Rat Brain Response to Chronic Rapamycin Treatment. In PLoS One on 2013 May 28 by Matylda Macias, Magdalena Blazejczyk, et al.. PMID: 23724051, , (2013)

[PMID:23724051](#)

el at., Tunneling Nanotubes Promote Intercellular Mitochondria Transfer Followed by Increased Invasiveness in Bladder Cancer Cells. In Oncotarget on 2017 Feb 28 by Jinjin Lu, Xiufen Zheng, et al.. PMID: 28107184, , (2017)

[PMID:28107184](#)

el at., Propranolol Sensitizes Thyroid Cancer Cells to Cytotoxic Effect of Vemurafenib. In Oncol Rep on 2016 Sep by Wei-Jun Wei, Chen-Tian Shen et

al..PMID:27432558, , (2016)

[PMID:27432558](#)

el at., TDRG1 regulates chemosensitivity of seminoma TCam-2 cells to cisplatin via PI3K/Akt/mTOR signaling pathway and mitochondria-mediated apoptotic pathway.In Cancer Biol Ther on 2016 Jul 2 by Yu Gan , Yong Wang et al..PMID:27104982, , (2016)

[PMID:27104982](#)

el at., Long-term Stress with Hyperglucocorticoidemia-induced Hepatic Steatosis with VLDL Overproduction Is Dependent on both 5-HT2 Receptor and 5-HT Synthesis in Liver.In Int J Biol Sci on 2016 Jan 1 by Jihua Fu , Shaoxin Ma et al..PMID: 26884719, , (2016)

[PMID:26884719](#)

el at., H2O2 treatment or serum deprivation induces autophagy and apoptosis in naked mole-rat skin fibroblasts by inhibiting the PI3K/Akt signaling pathway.In Oncotarget on 2016 Dec 20 by Shanmin Zhao, Li Li et al..PMID:27863375, , (2016)

[PMID:27863375](#)

el at., Effective treatment with combination of peripheral 5 α - α -hydroxytryptamine synthetic inhibitor and 5 α - α -hydroxytryptamine 2 receptor antagonist on glucocorticoid α - α -induced whole α - α -body insulin resistance with hyperglycemia.In J Diabetes Investig on 2016 Nov by Shaoxin Ma , Tao Li et al..PMID:27177506, , (2016)

[PMID:27177506](#)

el at., Amlexanox Enhances Temozolomide-Induced Antitumor Effects in Human Glioblastoma Cells by Inhibiting IKBKE and the Akt-mTOR Signaling Pathway. In ACS Omega on 2021 Feb 5 by Jinbiao Xiong, Gaochao Guo, et al..PMID:33644550, , (2020)

[PMID:33644550](#)

el at., A Novel Late-Stage Autophagy Inhibitor That Efficiently Targets Lysosomes Inducing Potent Cytotoxic and Sensitizing Effects in Lung Cancer. In Cancers (Basel) on 2022 Jul 12 by Adri α Molero-Valenzuela, Pere Fontova, et al..PMID: 35884450, , (2022)

[PMID:35884450](#)

el at., The AMPK-dependent inhibition of autophagy plays a crucial role in protecting photoreceptor from photooxidative injury In J Photochem Photobiol B On2023 AugbyYu-Lin Li , Tian-Zi Zhang et al..PMID:37302163, , (2023)

[PMID:37302160](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.