SHP-2(Phospho-Tyr542) Antibody

Catalog No: #11319

Package Size: #11319-1 50ul #11319-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	SHP-2(Phospho-Tyr542) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of SHP-2 only when phosphorylated at tyrosine 542.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 542 (H-E-Y(p)-T-N) derived from Human SHP-2.
•	
Conjugates	Unconjugated
	Unconjugated SHP-2
Conjugates	, s
Conjugates Target Name	SHP-2
Conjugates Target Name Modification	SHP-2 Phospho
Conjugates Target Name Modification Other Names	SHP-2 Phospho PTN11; PTP-1D; PTP-2C; PTP2C; PTPN11
Conjugates Target Name Modification Other Names Accession No.	SHP-2 Phospho PTN11; PTP-1D; PTP-2C; PTP2C; PTPN11 Swiss-Prot: Q06124NCBI Protein: NP_002825.3
Conjugates Target Name Modification Other Names Accession No. Concentration	SHP-2 Phospho PTN11; PTP-1D; PTP-2C; PTP2C; PTPN11 Swiss-Prot: Q06124NCBI Protein: NP_002825.3 1.0mg/ml
Conjugates Target Name Modification Other Names Accession No. Concentration	SHP-2 Phospho PTN11; PTP-1D; PTP-2C; PTP2C; PTPN11 Swiss-Prot: Q06124NCBI Protein: NP_002825.3 1.0mg/ml Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%


Application Details

Predicted MW: 72kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from 293 cells untreated or treated with starvation using SHP-2(Phospho-Tyr542) Antibody #11319.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using SHP-2(Phospho-Tyr542) Antibody #11319(left) or the same antibody preincubated with blocking peptide(right).

Background

Acts downstream of various receptor and cytoplasmic protein tyrosine kinases to participate in the signal transduction from the cell surface to the nucleus.

Ferjoux G, et al. (2003) Mol Biol Cell. 2003; 14(9): 3911-3928.

Shi ZQ, et al. (2000) Mol Cell Biol. 20(5): 1526-1536.

Li C, Friedman JM. (1999) Proc Natl Acad Sci U S A; 96(17): 9677-9682

Manes S, et al. (1999) Mol Cell Biol. 19(4): 3125-3135. Oh ES, et al. (1999) Mol Cell Biol; 19(4): 3205-3215.

Published Papers

el at., Glutamate Acting on N-Methyl-D-aspartate Receptors Attenuates Insulin-like Growth Factor-1 Receptor Tyrosine Phosphorylation and Its Survival Signaling Properties in Rat Hippocampal Neurons. In J Biol Chem 2009 Jan 9 by Wen-Hua Zheng, Rθ mi Quirioni ε t al..PMID: 18981172, , (2009)

PMID:18981172

el at., Glutamate acting on NMDA receptors attenuates IGF-1 receptor tyrosine phosphorylation and its survival signaling properties in rat hippocampal neurons. In J Biol Chem

on 2009 Jan 9 by Wen-Hua Zheng, R θ mi Quirioni ?et al..PMID:18981172, , (2009)

PMID:18981172

Note: This product is for in vitro research use only and is not intended for use in humans or animals.