

PI3 Kinase p85/p55 (phospho-Tyr467/199)Antibody

Catalog No: #11508

Package Size: #11508-1 50ul #11508-2 100ul

Description

Product Name	PI3 Kinase p85/p55 (phospho-Tyr467/199)Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB,IF
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total PI3 Kinase p85/p55 only when phosphorylated at tyrosine 467/199.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of tyrosine 467 (L-Y(p)-E-E-Y) derived from Human PI3 Kinase p85/p55.
Conjugates	Unconjugated
Target Name	PI3 Kinase p85/p55
Modification	Phospho
Other Names	p85, AGM7, p85-ALPHA, p55, p55-GAMMA, PIK3R1
Accession No.	Swiss-Prot:P27986Gene ID:5295
SDS-PAGE MW	55,85kd
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


Application Details

Predicted MW: 55, 85kd


Western blotting: 1:500~1:1000

IF 1:100-1:500

Images



Western blot analysis of extracts from 293 cells untreated or treated with sorbitol using PI3 Kinase p85/p55 (phospho-Tyr467/199)Antibody #11508.



All lanes : PI3 Kinase p85/p55 (phospho-Tyr467/199)
Antibody at 1/500 dilution Lane 1 : MCF-7 whole cell lysates
Lane 2 : MCF-7 treated with 100ng/mL EGF for 15 min whole cell lysates Lane 3 : MCF-7 treated with 200nM Dasatinib for 24hr whole cell lysates Lane 4 : JK whole cell lysates Lane 5 : JK treated with 2mM H2O2 for 15 min whole cell lysates
Lysates/proteins at 20 µg per lane. Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
Predicted band size: 55 kDa Observed band size: 60,85 kDa
Exposure time: 9 seconds

Background

Binds to activated (phosphorylated) protein-Tyr kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane. Necessary for the insulin-stimulated increase in glucose uptake and glycogen synthesis in insulin-sensitive tissues. Binds to activated (phosphorylated) protein-tyrosine kinases through its SH2 domain and regulates their kinase activity. During insulin stimulation, it also binds to IRS-1./Binds to activated (phosphorylated) protein-tyrosine kinases, through its SH2 domain, and acts as an adapter, mediating the association of the p110 catalytic unit to the plasma membrane.

Stoyanov, B. et al. (1995) Science 269, 690-3.

Simpson, L. and Parsons, R. (2001) Exp Cell Res 264, 29-41.

Neri, L.M. et al. (2002) Biochim Biophys Acta 1584, 73-80.

Published Papers

el at., Cardioprotection of (%2B)-sodium 5-bromo-2-(α -hydroxypentyl) benzoate (BZP) on mouse myocardium I/R injury through inhibiting 12/15-LOX-2 activity. In J Mol Cell Cardiol on 2019 Oct by Xiao Y, Song C, et al..PMID:31362020, , (2019)

[PMID:31362020](#)

el at., Okadaic acid promotes epithelial mesenchymal transition of hepatocellular carcinoma cells by inhibiting protein phosphatase 2A.In J Cell Biochem on 2020 Jan 6. by Kong J, Li D, et al..PMID: 31904141 , , (2020)

[PMID:31587272](#)

el at., Isoliquiritigenin inhibits the proliferation, migration and metastasis of Hep3B cells via suppressing cyclin D1 and PI3K/AKT pathway.In Biosci Rep on 2020 Jan 31; by Huang Y, Liu C, et al..PMID: 31840737, , (2020)

[PMID:31840737](#)

el at., Positive feedback loop of FAM83A/PI3K/AKT/c-Jun induces migration, invasion and metastasis in hepatocellular carcinoma.In Biomed Pharmacother on 2020 Mar; by Liu C, Peng X et al..PMID: 31901550, , (2020)

[PMID:31901550](#)

el at., Combination of Ruthenium Complex and Doxorubicin Synergistically Inhibits Cancer Cell Growth by Down-Regulating PI3K/AKT Signaling Pathway.In Front Oncol on 2020 Feb 18 by Lin K, Rong Y, et al..PMID:32133289, , (2020)

[PMID:32133289](#)

el at., Dexmedetomidine protects against lipopolysaccharide-induced acute kidney injury by enhancing autophagy through inhibition of the PI3K/AKT/mTOR pathway. In Front Pharmacol. On 2020 Feb 25; by Zhao Y, Feng X, et al..PMID: 32158395, , (2020)

[PMID:32158395](#)

el at., Angiopoietin-like 3 induces podocyte F-actin rearrangement through integrin $\alpha_5\beta_1$ FAK/PI3K pathway-mediated Rac1 activation. In Biomed Res Int on 2013 by Lin Y, Rao J, et al..PMID:24294595, , (2013)

[PMID:24294595](#)

el at., Angiopoietin-Like 3 Induces Podocyte F-Actin Rearrangement through Integrin /FAK/PI3K Pathway-Mediated Rac1 Activation. In Biomed Res Int on 2013 by Yi Lin, Jia Rao, et al..PMID: 24294595, , (2013)

[PMID:24294595](#)

el at., An alternatively spliced variant of CXCR3 mediates the metastasis of CD133+ liver cancer cells induced by CXCL9. In Oncotarget on 2016 Mar 22 by Qiang Ding, Yujia Xia et al..PMID:26883105, , (2016)

[PMID:26883105](#)

el at., A Novel Long Non-Coding RNA lnc030 Maintains Breast Cancer Stem Cell Stemness by Stabilizing SQLE mRNA and Increasing Cholesterol Synthesis. In Adv Sci (Weinh) On 2020 Nov 30 by Yilu Qin 1, Yixuan Hou, Shuiqing Liu, et al., , (2020)

[PMID:33511005](#)

el at., A Novel Long Non-Coding RNA lnc030 Maintains Breast Cancer Stem Cell Stemness by Stabilizing SQLE mRNA and Increasing Cholesterol Synthesis. In Adv Sci (Weinh) on 2020 Nov 30 by Yilu Qin, Yixuan Hou, et al..PMID:33511005, , (2021)

[PMID:33511005](#)

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF- κ B Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy. In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang, et al..PMID: 35370667, , (2022)

[PMID:35370667](#)

el at., Monascus vinegar alleviates high-fat-diet-induced inflammation in rats by regulating the NF- κ B and PI3K/AKT/mTOR pathways, , (2022)

[PMID:](#)

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF- κ B Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy. In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang, et al..PMID: 35370667, , (2022)

[PMID:35370667](#)

el at., Alleviative effects of α -lipoic acid on muscle atrophy via the modulation of TNF- α /JNK and PI3K/AKT pathways in high-fat diet and streptozotocin-induced type 2 diabetic rats. In Food Sci Nutr On 2023 Jan 12 by Chih-Yuan Ko 1 2 3, Chi-Hao Wu et al..PMID: 37051351, , (2023)

[PMID:37051351](#)

el at., Histone Methyltransferase KMT2B Promotes Metastasis and Angiogenesis of Cervical Cancer by Upregulating EGF Expression. In Int J Biol Sci On 2023 Jan 15 by Dan Zhao, Hui Yuan et al..PMID:36594087, , (2023)

[PMID:36594087](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.