#### **Product Datasheet**

# Cyclin E1(phospho-Thr395) Antibody

Catalog No: #11541

Package Size: #11541-1 50ul #11541-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_		
11)6	29Cr	ıntınn
$\boldsymbol{\nu}$		iption

Product Name	Cyclin E1(phospho-Thr395) Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.	
	Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho	
	specific antibodies were removed by chromatogramphy using non-phosphopeptide.	
Applications	WB	
Species Reactivity	Hu	
Specificity	The antibody detects endogenous level of Cyclin E1 only when phosphorylated at threonine 395.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around phosphorylation site of threonine 395 (L-L-T(p)-P-P)derived from Human Cyclin E1	
Conjugates	Unconjugated	
Target Name	Cyclin E1	
Modification	Phospho	
Other Names	CCNE; CCNE1;	
Accession No.	Swiss-Prot: P24864NCBI Protein: NP_001229.1	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

## **Application Details**

Predicted MW: 48kd
Western blotting: 1:1000

#### **Images**



Western blot analysis of extracts from 293 cells untreated or treated with hydroxyurea using Cyclin E1(phospho-Thr395) Antibody #11541.

## Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK2, whose activity is required for cell cycle G1/S transition.

Won K.A., Reed S.I.EMBO J. 15:4182-4193(1996)

Welcker M., Singer J., Loeb K.R., Grim J., Bloecher A., Mol. Cell 12:381-392(2003)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.