MEK1 (Phospho-T292) Rabbit mAb

Catalog No: #13408

Package Size: #13408-1 50ul #13408-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| | | 4.5 | |
|--------------------|-------|------|-----|
| | escri | ntir | าท |
| $\boldsymbol{\nu}$ | COUL | μu | ווע |
| | | | |

| • | |
|-----------------------|--|
| Product Name | MEK1 (Phospho-T292) Rabbit mAb |
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Clone No. | SD2088 |
| Purification | ProA affinity purified |
| Applications | WB |
| Species Reactivity | Hu |
| Immunogen Description | Synthetic phospho-peptide corresponding to residues surrounding Thr292 of human MEK1 |
| Conjugates | Unconjugated |
| Other Names | Dual specificity mitogen activated protein kinase kinase 1 antibody Dual specificity mitogen-activated protein |
| | kinase kinase 1 antibody ERK activator kinase 1 antibody MAP kinase kinase 1 antibody MAP kinase/Erk |
| | kinase 1 antibody MAP2K1 antibody MAPK/ERK kinase 1 antibody MAPKK 1 antibody MAPKK1 antibody |
| | MEK 1 antibody Mek1 antibody MEKK1 antibody Mitogen activated protein kinase kinase 1 antibody MKK 1 |
| | antibody MKK1 antibody MP2K1_HUMAN antibody PRKMK1 antibody Protein kinase mitogen activated |
| | kinase 1 (MAP kinase kinase 1) antibody Protein kinase mitogen activated, kinase 1 antibody |
| Accession No. | Swiss-Prot#:Q02750 |
| Calculated MW | Predicted band size: 43 kDa |
| SDS-PAGE MW | Observed band size: 43 kDa |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |
| | |

Application Details

WB: 1:500-1:2000

Images



All lanes: MEK1 (Phospho-T292) Rabbit mAb at 1/1k dilution

Lane 1: Hela whole cell lysates

Lane 2 : Hela treated with 100ng/mL Nocodazole for 17 hours whole cell lysates

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 43 kDa Observed band size: 43 kDa

Exposure time: 10 seconds

Background

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK 5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.