

GluR1(Phospho-S845) Rabbit mAb

Catalog No: #13418

Package Size: #13418-1 50ul #13418-2 100ul


Description

Product Name	GluR1(Phospho-S845) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	JJ2009
Purification	ProA affinity purified
Applications	WB
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser845 of human GluR1.
Conjugates	Unconjugated
Other Names	GLUR 1 antibody GLUR A antibody AMPA 1 antibody AMPA selective glutamate receptor 1 antibody AMPA-selective glutamate receptor 1 antibody GluA1 antibody GLUH1 antibody GluR K1 antibody GluR-1 antibody GluR-A antibody GluR-K1 antibody GLUR1 antibody GLURA antibody GluRK1 antibody Glutamate receptor 1 antibody Glutamate receptor ionotropic AMPA 1 antibody Glutamate receptor ionotropic antibody Glutamate receptor, ionotropic, AMPA 1 antibody Gria1 antibody GRIA1_HUMAN antibody HBGR1 antibody MGC133252 antibody OTTHUMP00000160643 antibody OTTHUMP00000165781 antibody OTTHUMP00000224241 antibody OTTHUMP00000224242 antibody OTTHUMP00000224243 antibody
Accession No.	Swiss-Prot#:P42261
Calculated MW	Predicted band size: 102 kDa
SDS-PAGE MW	Observed band size: 102 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:2000

Images



All lanes : GluR1(Phospho-S845) Rabbit mAb at 1/500 dilution

Lane 1 : Mouse brain tissue lysates

Lane 2 : Mouse brain tissue lysates treated with λpp for 1 hour

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 102 kDa

Observed band size: 102 kDa

Exposure time: 15 seconds

Background

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neuro-transmission by glutamate whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca^{2+} ions. The NMDA receptors consist of five subunits: epsilon 1, 2, 3, 4 and one zeta subunit. The zeta subunit is expressed throughout the brainstem whereas the four epsilon subunits display limited distribution.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.