#### **Product Datasheet**

# alpha Synuclein (Phospho-Ser129) Rabbit mAb

Catalog No: #13434

Package Size: #13434-1 50ul #13434-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


D	es	cri	pı	(IO	n
_					

Description	
Product Name	alpha Synuclein (Phospho-Ser129) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	JB22-44
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Ser129 of human alpha Synuclein.
Conjugates	Unconjugated
Other Names	Alpha synuclein antibody Alpha-synuclein antibody Alpha-synuclein, isoform NACP140 antibody alphaSYN
	antibody MGC105443 antibody MGC110988 antibody MGC127560 antibody MGC64356 antibody NACP
	antibody Non A beta component of AD amyloid antibody Non A4 component of amyloid antibody Non A4
	component of amyloid precursor antibody Non-A beta component of AD amyloid antibody Non-A-beta
	component of alzheimers disease amyloid , precursor of antibody Non-A4 component of amyloid precursor
	antibody Non-A4 component of amyloid, precursor of antibody OTTHUMP00000218549 antibody
	OTTHUMP00000218551 antibody OTTHUMP00000218552 antibody OTTHUMP00000218553 antibody
	OTTHUMP00000218554 antibody PARK 1 antibody PARK 4 antibody PARK1 antibody PARK4 antibody
	Parkinson disease (autosomal dominant, Lewy body) 4 antibody Parkinson disease familial 1 antibody SNCA
	antibody Snca synuclein, alpha (non A4 component of amyloid precursor) antibody SYN antibody Synuclein
	alpha antibody Synuclein alpha 140 antibody Synuclein, alpha (non A4 component of amyloid precursor)
	antibody SYUA_HUMAN antibody
Accession No.	Swiss-Prot#:P37840
Calculated MW	Predicted band size: 14 kDa
SDS-PAGE MW	Observed band size: 18 kDa
Formulation	Rabbit IgG in 10mM phosphate buffered saline , pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium
	azide and 50% glycerol.
Storage	Store at -20°C


### **Application Details**

WB: 1:500-1:2000 ICC/IF: 1:50-1:200 IHC: 1:50-1:200


## **Images**




All lanes: alpha Synuclein (Phospho-Ser129) Rabbit mAb at 1/1k dilutionLane 1: Human fetal Brain Iysates Lane 2: Mouse Brain IysatesLane 3: Mouse Brain Iysates treated with Lambda Protein Phosphatase for 30min Lane 4: Rat Brain IysatesLane 5: Rat Brain Iysates treated with Lambda Protein Phosphatase for 30min Lysates/proteins at 20 µg per lane.SecondaryAll lanes: Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilutionPredicted band size: 14 kDa Observed band size: 18 kDaExposure time: 9 seconds



Formalin-fixed, paraffin-embedded human brain tissue stained for alpha Synuclein (Phospho-S129) using 13434 at 1/100 dilution in immunohistochemical analysis. The picture on the right is blocked with the phospho peptide.



Immunocytochemistry/ Immunofluorescence alpha Synuclein (Phospho-S129) antibody (13434)
ICC/IF staining of alpha Synuclein (Phospho-S129)in HUVEC cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.
Samples were incubated with 13434 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.
Nuclei were counterstained with DAPI.



Immunocytochemistry/ Immunofluorescence alpha Synuclein (Phospho-S129) antibody (13434)
ICC/IF staining of alpha Synuclein (Phospho-S129)in
SH-SY-5Y cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.
Samples were incubated with 13434 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.
Nuclei were counterstained with DAPI.

#### Background

The synuclein family members, including  $\alpha$ -synuclein (also designated NACP for non- $\beta$ -Amyloid component) and  $\beta$ -synuclein, are predominantly expressed in the brain and are speculated to be involved in synaptic regulation and neuronal plasticity.  $\alpha$ -synuclein is localized to neuronal cell bodies and synapses.  $\alpha$ -synuclein was first identified as a component of Alzheimer's disease amyloid plaques. Abnormal platelet function in Alzheimer's disease has been demonstrated. During megakaryocytic differentiation  $\alpha$ -synuclein has been found to be upregulated, while  $\beta$ -synuclein is downregulated, indicating that coordinate expression of synucleins may be important during hematopoetic cell differentiation. A mutant form of  $\alpha$ -synuclein has been found in patients with early onset Parkinson's disease.

	r۵		

Note: This product is for in vitro research use only and is not intended for use in humans or animals.