

PLC γ 1 (phospho Tyr1253) Polyclonal Antibody

Catalog No: #13607

Package Size: #13607-1 50ul #13607-2 100ul

Description

Product Name	PLC γ 1 (phospho Tyr1253) Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Applications	WB,ELISA
Species Reactivity	Human,Mouse,Rat
Specificity	Phospho-PLC γ 1 (Y1253) Polyclonal Antibody detects endogenous levels of PLC γ 1 protein only when phosphorylated at Y1253.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human PLCG1 around the phosphorylation site of Tyr1253. AA range:1221-1270
Conjugates	Unconjugated
Other Names	PLCG1; PLC1; 1-phosphatidylinositol 4; 5-bisphosphate phosphodiesterase gamma-1; PLC-148; Phosphoinositide phospholipase C-gamma-1; Phospholipase C-II; PLC-II; Phospholipase C-gamma-1; PLC-gamma-1
Accession No.	Swiss Prot:P19174GeneID:5335
SDS-PAGE MW	150
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Background

phospholipase C gamma 1(PLCG1) Homo sapiens The protein encoded by this gene catalyzes the formation of inositol 1,4,5-trisphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. This reaction uses calcium as a cofactor and plays an important role in the intracellular transduction of receptor-mediated tyrosine kinase activators. For example, when activated by SRC, the encoded protein causes the Ras guanine nucleotide exchange factor RasGRP1 to translocate to the Golgi, where it activates Ras. Also, this protein has been shown to be a major substrate for heparin-binding growth factor 1 (acidic fibroblast growth factor)-activated tyrosine kinase. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],

Note: This product is for in vitro research use only and is not intended for use in humans or animals.