## **Product Datasheet**

## NF-κB2 p100 (phospho-Ser866/870) rabbit pAb

WB

Catalog No: #13673

Description

Applications

Package Size: #13673-1 50ul #13673-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| · · · · · · · · · · · · · · · · · · · |   |
|---------------------------------------|---|
| Product Name                          | NF-κB2 p100 (phospho-Ser866/870) rabbit pAb   |
| Host Species                          | Rabbit  |
| Clonality                             | Polyclonal  |
| Purification                          | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |

Species Reactivity Human, Mouse
Specificity This antibody detects endogenous levels of Human Mouse NF-kB2 p100 (phospho-Ser866 or 870)

Immunogen Description Synthesized phosho peptide around human NF-κB2 p100 (Ser866 and 870)
Conjugates Unconjugated

Other Names

Nuclear factor NF-kappa-B p100 subunit (DNA-binding factor KBF2) (H2TF1) (Lymphocyte translocation chromosome 10 protein) (Nuclear factor of kappa light polypeptide gene enhancer in B-cells 2) (Oncogene

Lyt-10) (Lyt10) [Cleaved into: Nuclear factor NF-kappa-B p52 subunit]

Accession No. Swiss Prot:Q00653GeneID:4791

SDS-PAGE MW 100

Concentration 1 mg/ml

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Storage -20°C/1

## **Application Details**

WB 1:1000-2000

## Background

nuclear factor kappa B subunit 2(NFKB2) Homo sapiens This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013],

Note: This product is for in vitro research use only and is not intended for use in humans or animals.