Product Datasheet

Mucin 1 (phospho Ser1227) Polyclonal Antibody

Catalog No: #13697

Package Size: #13697-1 50ul #13697-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	Mucin 1 (phospho Ser1227) Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific
	immunogen.
Applications	IHC;IF;ELISA
Species Reactivity	Human
Specificity	Phospho-Mucin 1 (S1227) Polyclonal Antibody detects endogenous levels of Mucin 1 protein only when
	phosphorylated at S1227.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human MUC1 around the
	phosphorylation site of Ser1227. AA range:1196-1245
Conjugates	Unconjugated
Other Names	MUC1; PUM; Mucin-1; MUC-1; Breast carcinoma-associated antigen DF3; Carcinoma-associated mucin;
	Episialin; H23AG; Krebs von den Lungen-6; KL-6; PEMT; Peanut-reactive urinary mucin; PUM; Polymorphic
	epithelial mucin; PEM; Tumor-associated ep
Accession No.	Swiss Prot:P15941GeneID:4582
Concentration	1 mg/ml

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Application Details

IHC 1:100-1:300; ELISA 1:5000; IF 1:50-200

Background

Formulation

Storage

mucin 1, cell surface associated(MUC1) Homo sapiens This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate sp

Note: This product is for in vitro research use only and is not intended for use in humans or animals.

-20°C/1