

CREB-2 (phospho Ser245) Polyclonal Antibody

Catalog No: #13948

Package Size: #13948-1 50ul #13948-2 100ul

Description

Product Name	CREB-2 (phospho Ser245) Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Applications	IHC-p,IF(paraffin section),ELISA
Species Reactivity	Human
Specificity	Phospho-CREB-2 (S245) Polyclonal Antibody detects endogenous levels of CREB-2 protein only when phosphorylated at S245.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human ATF4 around the phosphorylation site of Ser245. AA range:212-261
Conjugates	Unconjugated
Other Names	ATF4; CREB2; TXREB; Cyclic AMP-dependent transcription factor ATF-4; cAMP-dependent transcription factor ATF-4; Activating transcription factor 4; Cyclic AMP-responsive element-binding protein 2; CREB-2; cAMP-responsive element-binding prot
Accession No.	Swiss Prot:P18848GeneID:468
Calculated MW	38kd
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

WB 1:500-2000 ,Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Background

activating transcription factor 4(ATF4) Homo sapiens This gene encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromosome at q28 in a region containing a large inverted duplication. [provid

Note: This product is for in vitro research use only and is not intended for use in humans or animals.