

c-Myc (phospho Thr358) Polyclonal Antibody

Catalog No: #13959

Package Size: #13959-1 50ul #13959-2 100ul

Description

Product Name	c-Myc (phospho Thr358) Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Applications	IHC-p,IF(paraffin section),IP,ELISA
Species Reactivity	Human,Mouse,Rat
Specificity	Phospho-c-Myc (T358) Polyclonal Antibody detects endogenous levels of c-Myc protein only when phosphorylated at T358.
Immunogen Description	The antiserum was produced against synthesized peptide derived from human Myc around the phosphorylation site of Thr358. AA range:325-374
Conjugates	Unconjugated
Other Names	MYC; BHLHE39; Myc proto-oncogene protein; Class E basic helix-loop-helix protein 39; bHLHe39; Proto-oncogene c-Myc; Transcription factor p64
Accession No.	Swiss Prot:P01106GeneID:4609
SDS-PAGE MW	50,
Concentration	1 mg/ml
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage	-20°C/1

Application Details

Immunohistochemistry: 1/100 - 1/300. Immunoprecipitation: 2-5 ug/mg lysate. ELISA: 1/20000. Not yet tested in other applications.

Background

v-myc avian myelocytomatosis viral oncogene homolog(MYC) Homo sapiens The protein encoded by this gene is a multifunctional, nuclear phosphoprotein that plays a role in cell cycle progression, apoptosis and cellular transformation. It functions as a transcription factor that regulates transcription of specific target genes. Mutations, overexpression, rearrangement and translocation of this gene have been associated with a variety of hematopoietic tumors, leukemias and lymphomas, including Burkitt lymphoma. There is evidence to show that alternative translation initiations from an upstream, in-frame non-AUG (CUG) and a downstream AUG start site result in the production of two isoforms with distinct N-termini. The synthesis of non-AUG initiated protein is suppressed in Burkitt's lymphomas, suggesting its importance in the normal function of this gene. [provided by RefSeq, Jul 2008],

Note: This product is for in vitro research use only and is not intended for use in humans or animals.