

STAT3 (Phospho-Ser727) Rabbit mAb

Catalog No: #14152

Package Size: #14152-1 50ul #14152-2 100ul



SAB[®]
Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com

Description

Product Name	STAT3 (Phospho-Ser727) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	SR2312
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB, ICC/IF, IHC
Species Reactivity	Human Mouse Rat
Specificity	Phospho-STAT3 (S727) Antibody detects endogenous levels of total Phospho-STAT3 (S727)
Immunogen Description	A synthesized peptide derived from human Phospho-STAT3 (S727)
Conjugates	Unconjugated
Other Names	APRF;Acute-phase response factor;Signal transducer and activator of transcription 3;Stat3;
Accession No.	Uniprot:P40763
Calculated MW	Predicted band size: 88 kDa
SDS-PAGE MW	Observed band size: 98 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4Λ C short term. Store at -20Λ C long term. Avoid freeze / thaw cycle.


Application Details

WB: 1:500-1:2000


ICC/IF: 1:50-1:200

IHC: 1:50-1:200


Images




All lanes : STAT3 (Phospho-Ser727) Rabbit mAb at 1/1k dilution
Lane 1 :A431 cell lysate Lane 2 :A431 cell lysate treated with 100ng/ml EGF 10min
lysate/proteins at 20 µg per lane.SecondaryAll lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
Predicted band size: 88 kDa Observed band size: 98 kDaExposure time: 5 seconds




Formalin-fixed, paraffin-embedded rat brain tissue stained for STAT3 (Phospho-Ser727) using 14152 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded rat kidney tissue stained for STAT3 (Phospho-Ser727) using 14152 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded human brain tissue stained for STAT3 (Phospho-Ser727) using 14152 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/Immunofluorescence STAT3 (Phospho-Ser727) antibody (14152)
 ICC/IF staining of STAT3 (Phospho-Ser727) in A431 cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with 14152 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500. Nuclei were counterstained with DAPI.

Published Papers

Aysegul Dalmizrak;Nur Selvi Gunel;Burcin Tezcanh Kaymaz;Fahri Sahin;Guray Saydam;Buket Kosova el at., Effects of Rituximab on JAK-STAT and NF- κ B signaling pathways in acute lymphoblastic leukemia and chronic lymphocytic leukemia, , (2019)

PMID:

Note: This product is for in vitro research use only and is not intended for use in humans or animals.