

EGFR (Phospho-Y1197) Rabbit mAb

Catalog No: #14180

Package Size: #14180-1 50ul #14180-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	EGFR (Phospho-Y1197) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;IF
Species Reactivity	Human
Specificity	Phospho-EGFR (Y1197) Antibody detects endogenous levels of Phospho-EGFR (Y1197)
Immunogen Description	A synthesized peptide derived from human EGFR around the phosphorylation site of Y1197
Conjugates	Unconjugated
Other Names	EC 2.7.10.1; ERBB1; Epidermal growth factor receptor precursor; kinase EGFR;
Accession No.	Uniprot:P00533
Calculated MW	134 kDa
SDS-PAGE MW	175 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4ℳ C short term. Store at -20ℳ C long term. Avoid freeze / thaw cycle.

Application Details


WB:1:500~1:2000

IHC:1:50~1:100

ICC/IF:1:50~1:100

IP:1:20

Images



All lanes : Phospho-EGFR (Y1197) Rabbit mAb at 1/1k dilution

Lane 1 : Hela whole cell lysates

Lane 2 : Hela treated with 100ng/ml EGF for 20min whole cell lysates


Lysates/proteins at 20 µg per lane.

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution


Predicted band size: 134 kDa

Observed band size: 175 kDa

Exposure time: 4 seconds



Formalin-fixed, paraffin-embedded human lung carcinoma tissue stained for Phospho-EGFR (Y1197) using 14180 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence - p-EGFR (Y1197) antibody (14180)

14180 staining p-EGFR (Y1197) in the HeLa cell line by ICC/IF. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 14180 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

The negative control is shown in bottom right hand panel - for the negative control, purified 14180 was used at a dilution of 1/100 followed by Alexa FluorB 647 goat anti- rabbit antibody at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.