

eIF4B (Phospho-Ser406) Rabbit mAb

Catalog No: #14278

Package Size: #14278-1 50ul #14278-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description


Product Name	eIF4B (Phospho-Ser406) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Isotype	Rabbit IgG
Purification	Affinity-chromatography
Applications	WB;IHC;IF
Species Reactivity	Human;Mouse;Rat
Specificity	Phospho-eIF4B (S406) Antibody detects endogenous levels of total Phospho-eIF4B (S406)
Immunogen Description	A synthesized peptide derived from human Phospho-eIF4B (S406)
Conjugates	Unconjugated
Other Names	EIF4B;
Accession No.	Uniprot:P23588
Calculated MW	69 kDa
SDS-PAGE MW	75 kDa
Formulation	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at +4 Λ C short term. Store at -20 Λ C long term. Avoid freeze / thaw cycle.

Application Details


WB:1:500~1:2000

IHC:1:50~1:200


Images




All lanes : Phospho-eIF4B (S406) Rabbit mAb at 1/1k dilution
 Lane 1 : Hela whole cell lysates
 Lane 2 : 293T whole cell lysates
 Lane 3 : K562 whole cell lysates
 Lane 4 : HepG2 whole cell lysates
 Lane 5 : 3T3 whole cell lysates
 Lysates/proteins at 20 μ g per lane.
 Secondary: Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 69 kDa
 Observed band size: 75 kDa
 Exposure time: 3 seconds




All lanes : Phospho-eIF4B (S406) Rabbit mAb at 1/1k dilution
 Lane 1 : Mouse Liver lysates
 Lane 2 : Rat Liver lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 69 kDa
 Observed band size: 75 kDa
 Exposure time: 3 seconds




All lanes : Phospho-eIF4B (S406) Rabbit mAb at 1/1k dilution
 Lane 1 : Hela whole cell lysates
 Lane 2 : Hela treated with 100nM Insulin for 20min whole cell lysates
 Lane 3 : 293T whole cell lysates
 Lane 4 : 293T treated with Lambda Protein Phosphatase for 30min whole cell lysates
 Lane 5 : 293T treated with 100nM Insulin for 20min whole cell lysates
 Lysates/proteins at 20 µg per lane.
 Secondary All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution
 Predicted band size: 69 kDa
 Observed band size: 75 kDa
 Exposure time: 6 seconds



Formalin-fixed, paraffin-embedded mouse lung tissue stained for Phospho-eIF4B (S406) using 14278 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded rat lung tissue stained for Phospho-eIF4B (S406) using 14278 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence p-eIF4B (S406) antibody (14278) 14278 staining p-eIF4B (S406) in the HeLa cell line by ICC/IF. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100. Samples were incubated with 14278 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500. The negative control is shown in bottom right hand panel - for the negative control, purified 14278 was used at a dilution of 1/100 followed by Alexa FluorB 647 goat anti- rabbit antibody at a dilution of 1/500. Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.