## Met (Phospho-Tyr1234/1235) Rabbit mAb

Catalog No: #14316

Package Size: #14316-1 50ul #14316-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


## Description

Product Name	Met (Phospho-Tyr1234/1235) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	SR4356
Purification	Protein A
Applications	WB, ICC/IF, IHC
Species Reactivity	Human;Mouse;Rat
Specificity	This antibody detects endogenous levels of Met only when phosphorylated at Tyr1234 or Thr1235,and dually
	phosphorylated at two sites.
Target Name	Met
Modification	Phospho
Other Names	Hepatocyte growth factor receptor;HGF receptor;HGF/SF receptor;Proto-oncogene c-Met;Scatter factor
	receptor;SF receptor;Tyrosine-protein kinase Met;
Accession No.	P08581
Calculated MW	Predicted band size: 155 kDa
SDS-PAGE MW	Observed band size: 155kDa
Formulation	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Storage	Store at -20°C /1 year


## **Application Details**

WB: 1:500-1:2000 ICC/IF: 1:50-1:200 IHC: 1:50-1:200


## **Images**



All lanes: Met (Phospho Tyr1234/1235) Rabbit mAb at 1/1k dilutionLane 1: Hela whole cell lysatesLane 2: Hela treated with 40 ng/ml HGF for 5 min whole cell lysatesLysates/proteins at 20 µg per lane. Secondary All lanes: Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution Predicted band size: 155 kDa Observed band size: 155kDa Exposure time: 5 seconds



Formalin-fixed, paraffin-embedded human lung cancer tissue stained for Met (Phospho-Tyr1234/1235) using 14316 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/Immunofluorescence Met (Phospho-Tyr1234/1235) antibody (14316) ICC/IF staining of Met (Phospho-Tyr1234/1235) in 293T cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 14316 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 647 goat anti rabbit, used at a dilution of 1/500.

The negative control is shown in bottom right hand panel - for the negative control.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.