

IκB-α(Ab-32/36) Antibody

Catalog No: #21122

Package Size: #21122-1 50ul #21122-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	IκB-α(Ab-32/36) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC IF
Species Reactivity	Hu Ms
Specificity	The antibody detects endogenous level of total IκB-α protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.30~34/34~38 (H-D-S-G-L- D-S -M-K) derived from Human IκB-α.
Conjugates	Unconjugated
Target Name	IκB-α
Other Names	I-kappa-B-alpha; IKBA; NF-kappaB inhibitor alpha; NFKBI; NFKBIA
Accession No.	Swiss-Prot: P25963NCBI Protein: NP_065390.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 39kd

Western blotting: 1:500~1:1000


Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from 293 and 3T3 cells using IκB-α(Ab-32/36) Antibody #21122.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using IkB-a(Ab-32/36) Antibody #21122(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed HeLa cells using IkB-a(Ab-32/36) Antibody #21122.

Background

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.

Mattioli I, et al. (2004) *J Immunol*; 172(10): 6336-44.

Courtois G, et al. (2003) *J Clin Invest*; 112(7): 1108-15.

Nair A, et al. (2003) *Oncogene*; 22(1): 50-8.

Fan C, et al. (2002) *J Cell Sci*; 115(Pt 24): 4843-53.

Schubert SY, et al. (2002) *FASEB J*; 16(14): 1931-3.

Published Papers

el at., Astragalus polysaccharides exerts anti-infective activity by inducing human cathelicidin antimicrobial peptide LL-37 in respiratory epithelial cells. *In Phytother Res*. On 2018 Aug by Zhao L, Tan S et al..PMID: 29672953, , (2018)

[PMID:29672953](#)

el at., Ajudecumin A from *Ajuga ovalifolia* var. *calantha* exhibits anti-inflammatory activity in lipopolysaccharide-activated RAW264.7 murine macrophages and animal models of acute inflammation. *In Pharm Biol*. On 2018 Dec by Zhang H, Ren QC et al..PMID: 31070535, , (2018)

[PMID:31070535](#)

el at., HMG2 regulates non-tuberculous mycobacteria survival via modulation of M1 macrophage polarization. *In J Cell Mol Med* on 2019 Dec by Wang X, Chen S, et al..PMID:31596045, , (2019)

[PMID:31596045](#)

el at., Inhibitory effects of eugenol on RANKL-induced osteoclast formation via attenuation of NF- κ B and MAPK pathways. *In Connect Tissue Res* on 2015 Jun by Vishwa Deepak, Abe Kasonga et al..PMID:25405641, , (2015)

[PMID:25405641](#)

el at., The Effects of Xiangqing Anodyne Spray on treating acute soft-Tissue injury mainly depend on suppressing activations of AKT and p38 pathways. *In Evid Based Complement Alternat Med* on 2016 by Shudong Wang, Tao Li et al..PMID:27190541, , (2016)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.