#### **Product Datasheet**

# 4E-BP1 (Ab-37) Antibody

Catalog No: #21215

Package Size: #21215-1 50ul #21215-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

The state of the s	
Product Name	4E-BP1 (Ab-37) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were
	purified by affinity-chromatography using epitope-specific peptide.
Applications	WB IHC
Species Reactivity	Hu Ms Rt
Specificity	The antibody detects endogenous level of total 4E-BP1 protein.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around aa.35~39 (S-T-T-P-G) derived from Human 4E-BP1.
Conjugates	Unconjugated
Target Name	4E-BP1
Other Names	EIF4EBP1; PHAS-1;
Accession No.	Swiss-Prot: Q13541NCBI Protein: NP_004086.1
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%
	sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.


### **Application Details**

Predicted MW: 18kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

### **Images**



Western blot analysis of extracts from HepG2 and 293 cells using 4E-BP1 (Ab-37) Antibody #21215.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using 4E-BP1 (Ab-37) Antibody #21215 (left) or the same antibody preincubated with blocking peptide (right).

# Background

4E-BP1 encodes one member of a family of translation repressor proteins. The protein directly interacts with eukaryotic translation initiation factor 4E (eIF4E), which is a limiting component of the multisubunit complex that recruits 40S ribosomal subunits to the 5' end of mRNAs. Interaction of this protein with eIF4E inhibits complex assembly and represses translation. This protein is phosphorylated in response to various signals including UV irradiation and insulin signaling, resulting in its dissociation from eIF4E and activation of mRNA translation.

Gingras AC, et al. Genes Dev Nov. 2005 Feb 15.

Mothe-Satney I, et al.(2000) J Biol Chem:15(21): 2852-64.

Gingras AC, et al. (1999) Genes Dev:13(11): 1422-37.

Lal L, et al.(2005) Blood;105(4): 1669-7.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.