Product Datasheet

SAPK/JNK(Ab-183) Antibody

Catalog No: #21241

Package Size: #21241-1 50ul #21241-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| | | 4.5 | |
|-----|-----|--------------------|------|
| Des | cri | ntic | ı'n |
| レしこ | 011 | ν u ϵ | /I I |
| | | | |

| Product Name | SAPK/JNK(Ab-183) Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were |
| | purified by affinity-chromatography using epitope-specific peptide. |
| Applications | WB IHC |
| Species Reactivity | Hu Ms Rt |
| Specificity | The antibody detects endogenous level of total SAPK/JNK protein. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around aa.181~185 (M-M-T-P-Y) derived from Human SAPK/JNK. |
| Conjugates | Unconjugated |
| Target Name | SAPK/JNK |
| Other Names | JNK2 |
| Accession No. | Swiss-Prot:P45984Gene ID:5601 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |
| | |


Application Details

Predicted MW: 46 54 kd
Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

Images



Western blot analysis of extracts from Hela and MCF cells using SAPK/JNK(Ab-183) Antibody #21241.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using SAPK/JNK(Ab-183) Antibody #21241(left) or the same antibody preincubated with blocking peptide(right).

Background

Responds to activation by environmental stress and pro-inflammatory cytokines by phosphorylating a number of transcription factors, primarily components of AP-1 such as c-Jun and ATF2 and thus regulates AP-1 transcriptional activity. In T-cells, JNK1 and JNK2 are required for polarized differentiation of T-helper cells into Th1 cells.

Ferrer, et al. (2003) Neuropathology & Applied Neurobiology 29: 23 Zhonghong Guan, et al. (1999) J Biol Chem, Vol. 274: 36200-36206 D.Margriet Ouwens1, et al. (2002)The EMBO Journal 21: 3782

Published Papers

Anastasia F. Thη— ε" venin, Chati L. Zony, Brian J. Bahnson el at., Activation by phosphorylation and purification of human c-Jun N-terminal kinase (JNK) isoforms in milligram amounts., Protein Expression and Purification, 75: 138η— C146(2011)

PMID:20709173

el at., Astragaloside IV Inhibits Isoprenaline ζ ½nduced Cardiac Fibrosis by Targeting the Reactive Oxygen Species/Mitogen ζ ³ctivated Protein Kinase Signaling Axis.In Mol Med Rep on 2017 Apr by Hongliang Dai ,Guizhi Jia ,et al..PMID:28260010, , (2017)

PMID:28260010

Note: This product is for in vitro research use only and is not intended for use in humans or animals.