Product Datasheet

Cyclin B1(Ab-147) Antibody

Catalog No: #21540

Package Size: #21540-1 50ul #21540-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | Cyclin B1(Ab-147) Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were |
| | purified by affinity-chromatography using epitope-specific peptide. |
| Applications | WB IHC IF |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of total Cyclin B1 protein. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around aa.145~149 (A-F-S-D-V) derived from Human Cyclin B1. |
| Conjugates | Unconjugated |
| Target Name | Cyclin B1 |
| Accession No. | Swiss-Prot: P14635NCBI Protein: NP_114172.1 |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% |
| | sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |


Application Details

Predicted MW: 60kd


Western blotting: 1:500~1:1000
Immunohistochemistry: 1:50~1:100

Immunofluorescence: 1:100~1:200


Images



Western blot analysis of extracts from HepG2 and HT29 cells using Cyclin B1(Ab-147) Antibody #21540.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Cyclin B1(Ab-147) Antibody #21540(left) or the same antibody preincubated with blocking peptide(right).



Immunofluorescence staining of methanol-fixed Hela cells using Cyclin B1(Ab-147) Antibody #21540.

Background

The protein encoded by Cyclin B1 is a regulatory protein involved in mitosis. The gene product complexes with p34(cdc2) to form the maturation-promoting factor (MPF). Two alternative transcripts have been found, a constitutively expressed transcript and a cell cycle-regulated transcript, that is expressed predominantly during G2/M phase. The different transcripts result from the use of alternate transcription initiation sites. Norbury, C. and Nurse, P. (1992) Annu. Rev. Biochem. 61, 441-470.

Atherton-Fessler, S. et al. (1993) Mol. Cell. Biol. 13, 1675-1685.

Galaktionov, K. et al. (1995) Genes Dev. 9, 1046-1058.

Published Papers

el at., PKM2 phosphorylates MLC2 and regulates cytokinesis of tumour cells.In Nat Commun on 2014 Nov 21 by Yuhui Jiang , Yugang Wang et al..PMID: 25412762, , (2014)

PMID:25412762

Note: This product is for in vitro research use only and is not intended for use in humans or animals.