#### **Product Datasheet**

# C-Fos Antibody

Catalog No: #21667

Package Size: #21667-1 50ul #21667-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_		4.6
	escri	ption
_	00011	Puon


Product Name	C-Fos Antibody	
Host Species	Rabbit	
Clonality	Polyclonal	
Purification	Antibodies were produced by immunizing rabbits with synthetic peptide and KLH conjugates. Antibodies were	
	purified by affinity-chromatography using epitope-specific peptide.	
Applications	WB	
Species Reactivity	Hu Ms Rt	
Specificity	The antibody detects endogenous level of total C-Fos protein.	
Immunogen Type	Peptide-KLH	
Immunogen Description	Peptide sequence around aa.28~32( S-Y-Y-H-S) derived from Human C-Fos.	
Conjugates	Unconjugated	
Target Name	C-Fos	
Other Names	p55; AP-1;	
Accession No.	Swiss-Prot: P01100NCBI Protein: NP_005243.1	
Concentration	1.0mg/ml	
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02%	
	sodium azide and 50% glycerol.	
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.	

### **Application Details**

Predicted MW: 62kd

Western blotting: 1:500~1:1000

# **Images**



Western blot analysis of extracts from A549, Hela and 3T3 cells using C-Fos Antibody #21667.

# Background

Nuclear phosphoprotein which forms a tight but non-covalently linked complex with the JUN/AP-1 transcription factor. In the heterodimer, FOS and JUN/AP-1 basic regions each seems to interact with symmetrical DNA half sites. On TGF-beta activation, forms a multimeric

SMAD3/SMAD4/JUN/FOS complex at the AP1/SMAD-binding site to regulate TGF-beta-mediated signaling. Has a critical function in regulating the Has a critical function in regulating the development of cells destined to form and maintain the skeleton. It is thought to have an important role in signal transduction, cell proliferation and differentiation.

Okazaki K., Sagata N.EMBO J. 14:5048-5059(1995)

Zhang Y., Feng X.H., Derynck R.Nature 394:909-913(1998)

Bossis G., Malnou C.E., Farras R. Mol. Cell. Biol. 25:6964-6979(2005)

### **Published Papers**

el at., Piperine alleviates osteoclast formation through the p38/cı  $\zeta$  os/NFATc1 signaling axis.In Biofactors on Nov-Dec 2015 by Vishwa Deepak , Marlena C Kruger et al..PMID: 26627060, , (2015)

PMID:26627060

Note: This product is for in vitro research use only and is not intended for use in humans or animals.