Product Datasheet

DGAT2 antibody

Catalog No: #22070

Package Size: #22070 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| Product Name | DGAT2 antibody |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Purified by antigen-affinity chromatography. |
| Applications | WB |
| Species Reactivity | Hu Ms Rt |
| Immunogen Description | A synthesized peptide derived from human DGAT2 |
| Conjugates | Unconjugated |
| Target Name | DGAT2 |
| Accession No. | NCBI Gene ID: 84649NCBI mRNA#: NM_032564NCBI Protein#: NP_115953 |
| Concentration | 1mg/ml |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |

Application Details

Predicted MW: 44kd

Western blotting: 1:1000-1:3000

Images



Western blot analysis DGAT2 using K562 whole cell lysates

Background

Acyl-CoA:diacylglycerol acyltransferase, or DGAT (EC 2.3.1.20), is responsible for the synthesis of triglycerides. It catalyzes a reaction in which diacylglycerol is covalently joined to long chain fatty acyl-CoAs.[supplied by OMIM]

| Note: This product is for in vitro research use only and is not intended for use in humans or animals. | | |
|--|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |