IL-33 Monoclonal Antibody

Catalog No: #26032

Package Size: #26032 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$		4.0
	escri	ption
	00011	Puon

IL-33 Monoclonal Antibody
Mouse
Monoclonal
mAb (Clone 12B3C4)
Immunoaffinity chromotography purified IgG
ELISA WB IHC
Hu Ms
Recombinant protein
Raised against the recombinant IL-33 protein.
Unconjugated
IL-33
IL-33 (12B3C4), Interleukin-33, Nuclear factor from high endothelial venules, NFHEV
Swiss-Prot:O95760Gene ID:90865
1mg/ml
Supplied in PBS containing 0.02% sodium azide.
Can be stored at -20°C, stable for one year.

Images



Western blot analysis of IL-33 using 125ng of recombinant IL-33 with IL-33 antibody at 1 ug/mL.



Immunohistochemistry of IL-33 in human lymph node tissue with IL-33 antibody at 5 ug/mL.

Background

Interleukin-33 (IL-33) is a recently identified member of the IL-1 family of cytokines whose other members include IL-1αβ, IL-1Ra and IL-18. Its receptor has been shown to be ST2, an IL-1 receptor family member that also acts as a negative regulator of TLR-IL-1R signaling and IL-1R accessory protein (IL-1RAcP). Receptor binding of IL-33 activates NF-κB and MAP kinases and induces the expression of TH2-associated cytokines such as IL-4, IL-5 and IL-6. Prolonged IL-33 treatment of mice led to the development of eosinophilia, splenomegaly, and severe pathological changes in mucosal organs such as lungs, esophagus and small intestine. Recent experiments have shown that IL-33 can also co-localize with heterochromatin and possesses transcriptional repressor activities, indicating that IL-33 may function as both a proinflammatory cytokine and an intracellular nuclear factor with transcriptional regulatory properties. Despite its predicted molecular weight, IL-33 will often run at higher molecular weight in SDS-PAGE.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.