

## NGFR Antibody

Catalog No: #32596

Package Size: #32596-1 50ul #32596-2 100ul


## Description

|                       |                                                                                                                                                              |
|-----------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name          | NGFR Antibody                                                                                                                                                |
| Host Species          | Rabbit                                                                                                                                                       |
| Clonality             | Polyclonal                                                                                                                                                   |
| Purification          | Antibodies were purified by affinity purification using immunogen.                                                                                           |
| Applications          | WB                                                                                                                                                           |
| Species Reactivity    | Human,Mouse,Rat                                                                                                                                              |
| Specificity           | The antibody detects endogenous level of total NGFR protein.                                                                                                 |
| Immunogen Type        | Peptide                                                                                                                                                      |
| Immunogen Description | A synthetic peptide of human NGFR.                                                                                                                           |
| Conjugates            | Unconjugated                                                                                                                                                 |
| Target Name           | NGFR                                                                                                                                                         |
| Other Names           | CD271; Gp80-LNGFR; TNFRSF16; p75(NTR); p75NTR                                                                                                                |
| Accession No.         | Swiss-Prot:P08138NCBI Gene ID:4804                                                                                                                           |
| SDS-PAGE MW           | 45KD                                                                                                                                                         |
| Concentration         | 1.0mg/ml                                                                                                                                                     |
| Formulation           | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage               | Store at -20°C                                                                                                                                               |

## Application Details

WB 1:500 - 1:2000

## Images



Western blot analysis of extracts of various cell lines, using NGFR at 1:1000 dilution.

## Background

The p75 neurotrophin receptor (p75NTR), a member of the TNF receptor superfamily, is distinguished by multiple cysteine-rich ligand-binding

domains, a single transmembrane sequence and a noncatalytic cytoplasmic domain (1). p75NTR displays paradoxical functions when acting alone or with other receptor proteins. Working in concert with Trk receptors, p75NTR recognizes neurotrophins and transmits trophic signals into the cell. Both p75NTR and TrkA are required to activate PI3K-Akt signaling, while TrkA can individually activate the MAP kinase pathway. In contrast, p75NTR, possibly through JNK, ensures appropriate apoptosis of injured neurons and improperly targeted neonatal neurons (2). The p75NTR protein undergoes sequential cleavage similar to APP and Notch. First,  $\alpha$ -secretase removes the p75NTR ectodomain, eliminating ligand-mediated signaling. At this point, the membrane-tethered cleavage product can still fine-tune Trk-mediated trophic actions.  $\gamma$ -secretase cleaves within the transmembrane domain to liberate the cytoplasmic tail from its membrane anchor and allow the p75NTR intracellular domain to translocate to the nucleus (3,4).

## Published Papers

el et al., Downregulation of KCTD12 contributes to melanoma stemness by modulating CD271. In Cancer Biol Med on 2019 Aug by Shen W, Li Y, et al.. PMID:31565480, , (2019)

[PMID:31565480](#)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.