

CDH11 Antibody

Catalog No: #48405

Package Size: #48405-1 50ul #48405-2 100ul



Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	CDH11 Antibody
Host Species	Mouse
Clonality	Monoclonal
Clone No.	A6-C12
Purification	ProA affinity purified
Applications	WB, ICC, IHC, FC
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Conjugates	Unconjugated
Other Names	CAD11 antibody CAD11_HUMAN antibody Cadherin 11 antibody Cadherin 11 type 2 antibody Cadherin 11 type 2 OB cadherin (osteoblast) antibody Cadherin-11 antibody Cdh11 antibody CDHOB antibody OB antibody OB-cadherin antibody OBcadherin antibody OSF-4 antibody OSF4 antibody Osteoblast cadherin antibody
Accession No.	Swiss-Prot#:P55287
Calculated MW	88 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:2,000IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of CDH11 on human CDH11 recombinant protein using anti-CDH11 antibody at 1/1,000 dilution.




Western blot analysis of CDH11 on HEK293 (1) and CDH11-hlgGfc transfected HEK293 (2) cell lysate using anti-CDH11 antibody at 1/1,000 dilution.




Western blot analysis of CDH11 on MCF-7 (1), Jurkat (2) and HEK293 (3) cell lysate using anti-CDH11 antibody at 1/1,000 dilution.



Immunohistochemical analysis of paraffin-embedded human bladder cancer tissue using anti-CDH11 antibody. Counter stained with hematoxylin.



ICC staining CDH11 (green) and Actin filaments (red) in HL-7702 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of HeLa cells with CDH11 antibody at 1/100 dilution (green) compared with an unlabelled control (cells without incubation with primary antibody; red).

Background

Cadherins are calcium dependent cell adhesion proteins, a family of Ca^{++} -dependent adhesion molecules that influence cell-cell binding and are critical to the maintenance of tissue structure and morphogenesis. OB-cadherin (osteoblast-cadherin, cadherin-11, OSF-4) has two forms, OB-cadherin-1 and OB-cadherin-2. OB-cadherin-2 has a truncated cytoplasmic domain, missing amino acids 694-796. Both OB-cadherins are expressed in osteoblastic cell lines with low expression seen in lungs, testis and brain. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.