Product Datasheet

Insulin Antibody

Catalog No: #48491

Package Size: #48491-1 50ul #48491-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | Insulin Antibody |
|-----------------------|---|
| Host Species | Mouse |
| Clonality | Monoclonal |
| Clone No. | A6-6 |
| Purification | ProA affinity purified |
| Applications | IHC |
| Species Reactivity | Hu, Ms, Rt |
| Immunogen Description | peptide |
| Conjugates | Unconjugated |
| Other Names | IDDM antibody IDDM1 antibody IDDM2 antibody ILPR antibody ins antibody INS_HUMAN antibody Insulin A |
| | chain antibody Insulin B chain antibody IRDN antibody MODY10 antibody Preproinsulin antibody Proinsulin |
| | antibody Proinsulin precursor antibody |
| Accession No. | Swiss-Prot#:P01308 |
| Calculated MW | 12 kDa |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |


Application Details

IHC: 1:100-1:1,000

Images



Immunohistochemical analysis of paraffin-embedded rat pancreas tissue using anti-Insulin antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human pancreas tissue using anti-Insulin antibody. Counter stained with hematoxylin.

Background

Insulin is a hormone with extensive effects on both metabolism and several other body systems. It causes most of the body's cells to take up glucose from the blood (including liver, muscle, and fat tissue cells), storing it as glycogen in the liver and muscle, and stops use of fat as an energy source. Insulin is synthesized as a precursor molecule, proinsulin, which is processed prior to its secretion. A- and B-peptides are joined together by a disulfide bond to form insulin, while the central portion of the precursor molecule is cleaved and released as the C-peptide.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.