Product Datasheet

NF-κB p65 Antibody

Catalog No: #48498

Package Size: #48498-1 50ul #48498-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


D	es	cri	pti	10	1

2 ccciiption	
Product Name	NF-κB p65 Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Peptide affinity purified
Applications	WB, IHC, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	This antibody is produced by immunizing rabbits with a synthetic peptide (KLH-coupled) corresponding to
	N-terminal NF-кВ p65.
Conjugates	Unconjugated
Other Names	Avian reticuloendotheliosis viral (v rel) oncogene homolog A antibody MGC131774 antibody NF kappa B
	p65delta3 antibody NFKB3 antibody Nuclear Factor NF Kappa B p65 Subunit antibody Nuclear factor
	NF-kappa-B p65 subunit antibody Nuclear factor of kappa light polypeptide gene enhancer in B cells 3
	antibody Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 antibody
	OTTHUMP00000233473 antibody OTTHUMP00000233474 antibody OTTHUMP00000233475 antibody
	OTTHUMP00000233476 antibody OTTHUMP00000233900 antibody p65 antibody p65 NF kappaB antibody
	p65 NFkB antibody relA antibody TF65_HUMAN antibody Transcription factor p65 antibody v rel avian
	reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B
	cells 3 (p65)) antibody V rel avian reticuloendotheliosis viral oncogene homolog A antibody v rel
	reticuloendotheliosis viral oncogene homolog A (avian) antibody V rel reticuloendotheliosis viral oncogene
	homolog A, nuclear factor of kappa light polypeptide gene enhancer in B cells 3, p65 antibody
Accession No.	Swiss-Prot#:Q04206
Calculated MW	65 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:100-1:200 FC: 1:50-1:100


Images




Western blot analysis of NF-κB p65 on different lysates using anti-NF-κB p65 antibody at 1/1000 dilution. Positive control: Lane1: Hela Lane2:A549 Lane3: PC12 Lane 4: Mouse embryonic stem cell Lane5: NIH/3T3



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-NF-kB p65 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human lung carcinoma tissue using anti-NF-o Ω^{1} 2o Ω^{1} 2B p65 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of Hela cells with NF-κB p65 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Goat anti rabbit IgG (FITC) was used as the secondary antibody.

Background

NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. In unstimulated cells, NF-kB is sequestered in the cytoplasm by lkB inhibitory proteins. NF-kB-activating agents can induce the phosphorylation of lkB proteins, targeting them for rapid degradation through the ubiquitin-proteasome pathway and releasing NF-kB to enter the nucleus where it regulates gene expression.

Published Papers

el at., Salinomycin promotes T-cell proliferation by inhibiting the expression and enzymatic activity of immunosuppressive indoleamine-2,3-dioxygenase in human breast cancer cells. In Toxicol Appl Pharmacol on 2020 Oct 1 by Yuwen Sheng, Zhonghui Zhang, et al..PMID:32822738, , (2020)

PMID:32822738

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF-KB Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy.In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang, et al..PMID: 35370667, , (2022)

PMID:35370667

el at., Resveratrol attenuates cigarette smoke extract induced cellular senescence in human airway epithelial cells by regulating the miR-34a/SIRT1/NF-kB pathway. In Medicine (Baltimore) on 2022 Nov 18 by Xiao-Li Zeng, Xin-Na Yang, et al..PMID: 36401446, , (2022)

PMID:36401446

el at., Effect of Fushengong Decoction on PTEN/PI3K/AKT/NF-kB Pathway in Rats With Chronic Renal Failure via Dual-Dimension Network Pharmacology Strategy. In Front Pharmacol on 2022 Mar 15 by Hongyu Luo, Munan Wang, et al..PMID: 35370667, (2022)

PMID:35370667

el at., A multi-omics approach based on 1H-NMR metabonomics combined with target protein analysis to reveal the mechanism of RIAISs on cervical carcinomaln Aging (Albany NY)On2022 Sep 27byChai Yanlan , Aibibai Jielili et al..PMID:36170024, , (2023)

PMID:36170024

Zhu Dan;Zheng Na;Deng Kebin;Li Liangchang el at., Aurantio-obtusin Alleviates Dry Eye Disease by Targeting NF-κB/NLRP3 Signaling in Rodent Models, , (2023)

PMID:

Qi Liu; Ziwei Liang; Jiapu Wang; Yuhui Wang; Jie Wang; Shaojie Wang; Zhi Du; Liqin Zhao; Yan Wei; Di Huang el at., Mannose-modified multifunctional iron-based nanozyme for hepatocellular carcinoma treatment by remodeling the tumor microenvironment., (2025)

PMID:39923382

Yanlan Chai Yanlan; Aibibai Jielili; Juan Wang; Haiying Tang; Zi Liu; Ping Wang el at., A multi-omics approach based on B'HH-NMR metabonomics combined with target protein analysis to reveal the mechanism of RIAISs on cervical carcinoma patients, (2023)

PMID:36170024

Rui Yang;Meng-Yan Deng;Lu-Kui Yang;Guan-Zhe Wang;Jun Ma;Qiang Wen;Na Gao;Hai-Ling Qiao el at., Identification of Cytochrome P450 2E1 as a Novel Target in Neuroinflammation and Development of Its Inhibitor Q11 as a Treatment Strategy., (2025)

PMID:40122152

Ang Li;Peng Liu;Jiaohong Gan;Weijun Fang;Anjie Liu el at., Phellodendrine Exerts Protective Effects on Intra-abdominal Sepsis by Inactivating AKT/NF-kB Signaling., , (2025)

PMID:39953352

Note: This product is for in vitro research use only and is not intended for use in humans or animals.