

CD11c Rabbit mAb

Catalog No: #48743

Package Size: #48743-1 50ul #48743-2 100ul


Description

Product Name	CD11c Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SI19-06
Purification	ProA affinity purified
Applications	WB, IHC, IP
Species Reactivity	Hu
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	95 alpha chain antibody 95 antibody CD 11c antibody CD11 antigen-like family member C antibody CD11c antibody CD11c antigen antibody Complement component 3 receptor 4 subunit antibody CR4 antibody Integrin alpha X antibody Integrin alpha X chain antibody Integrin alpha-X antibody Integrin aX antibody Integrin subunit alpha X antibody integrin, alpha X (antigen CD11C (p150), alpha polypeptide) antibody integrin, alpha X (complement component 3 receptor 4 subunit antibody ITAX_HUMAN antibody ITGAX antibody LEU M5 alpha subunit antibody Leu M5 antibody Leukocyte adhesion glycoprotein p150 95 alpha chain antibody Leukocyte adhesion glycoprotein p150 antibody Leukocyte adhesion receptor p150 95 antibody Leukocyte adhesion receptor p150 antibody Leukocyte surface antigen p150 95 alpha subunit antibody Leukocyte surface antigen p150 alpha subunit antibody Myeloid membrane antigen alpha subunit antibody p150 95 integrin alpha chain antibody p150 antibody p150/95 antibody SLEB6 antibody
Accession No.	Swiss-Prot#:P20702
Calculated MW	128 kDa(Predicted)
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200

Images



Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-CD11c antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD11c antibody. Counter stained with hematoxylin.

Background

Integrin α X (CD11C, leukocyte surface antigen p150,95, CR4, Axb2) is a type 1 transmembrane protein that traditionally combines with β 2 chain to form a leukocyte-specific integrin known as inactivated-C3b (iC3b) receptor 4 (CR4). Integrin α X/ β 2 shares similar properties of the α M/ β 2 integrin in mediating adherence of neutrophils and monocytes to stimulated endothelial cells, and in phagocytosis of complement coated particles. Abnormal expression of Integrin α X is characteristic of hairy cell leukemia (HCL) and is dependent upon activation of proto-oncogenes Ras and JunD. Proteins and DNA elements that influence transcription of Integrin α X include Sp1 and Sp1-like factors, AP-1 family, C/EBP, Oct-2 and PU.1. Integrin α X is present on monocyte derivative dendritic cells (DCs), macrophages and NK cells. Upon activation, DCs present in skin (Langerhans cells), lining of nose, lung, stomach, intestine and blood can migrate to lymphoid tissues and interact with T and B cells to initiate and shape the immune response.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.