

Smad5 Rabbit mAb

Catalog No: #48746

Package Size: #48746-1 50ul #48746-2 100ul



Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Smad5 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SI17-06
Purification	ProA affinity purified
Applications	WB, ICC, IHC, FC
Species Reactivity	Hu, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	DKFZp781C1895 antibody DKFZp781O1323 antibody Dwfc antibody hSmad5 antibody JV5 1 antibody JV5-1 antibody MAD homolog 5 antibody MAD, mothers against decapentaplegic homolog 5 antibody MADH 5 antibody MADH5 antibody Mothers against decapentaplegic homolog 5 antibody mothers against decapentaplegic, drosophila, homolog of, 5 antibody Mothers against DPP homolog 5 antibody MusMLP antibody SMA and MAD related protein 5 antibody SMAD 5 antibody SMAD family member 5 antibody SMAD, mothers against DPP homolog 5 antibody Smad5 antibody SMAD5_HUMAN antibody
Accession No.	Swiss-Prot#:Q99717
Calculated MW	52 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of Smad5 on PC-3M cell lysates using anti-Smad5 antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human lung tissue using anti-Smad5 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse lung tissue using anti-Smad5 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue using anti-Smad5 antibody. Counter stained with hematoxylin.




ICC staining Smad5 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Smad5 in A549 cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Smad5 in PC-3M cells (green). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Hela cells with Smad5 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Smad proteins, the mammalian homologs of the *Drosophila* Mothers against dpp (Mad) have been implicated as downstream effectors of TGF β /BMP signaling. Smad1 (also designated Madr1 or JV4-1), Smad5 and mammalian Smad8 (also designated Smad9 or MADH6) are effectors of BMP2 and BMP4 function while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGF β and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGF β signaling by interfering with TGF β -mediated phosphorylation of other Smad family members.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.