

Paxillin Rabbit mAb

Catalog No: #48774

Package Size: #48774-1 50ul #48774-2 100ul


Description

Product Name	Paxillin Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SY23-02
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	FLJ16691 antibody FLJ23042 antibody Paired box protein Pax 1 antibody PAX 1 antibody PAX1 antibody PAXI_HUMAN antibody Paxillin alpha antibody Paxillin antibody PXN antibody PXN protein antibody
Accession No.	Swiss-Prot#:P49023
Calculated MW	65 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200


Images




Western blot analysis of Paxillin on NIH/3T3 cells lysates using anti-Paxillin antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-Paxillin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue using anti-Paxillin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-Paxillin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse ovary tissue using anti-Paxillin antibody. Counter stained with hematoxylin.



ICC staining Paxillin in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Paxillin in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Paxillin in SKOV-3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Paxillin is a focal adhesion phosphoprotein that is localized to the cytoskeleton. Phosphorylation of paxillin has been shown to occur in response to PDGF treatment, v-Src transformation or cross-linking of integrins. FAK (focal adhesion kinase) and PYK2 have been shown to phosphorylate paxillin. FAK phosphorylates paxillin specifically on Tyr 118 in vitro. However, FAK phosphorylation does not seem to be required for the recruitment of paxillin to cell adhesion sites. Paxillin may play a role in signal transduction, regulation of cell morphology and the recruitment of structural and signaling molecules to focal adhesions. It has been shown that the amount of paxillin is reduced in mitotic cells by proteolytic downregulation and that paxillin is alternatively phosphorylated on serine rather than on tyrosine and serine during mitosis.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.