

c-Jun Rabbit mAb

Catalog No: #48820

Package Size: #48820-1 50ul #48820-2 100ul

Description

Product Name	c-Jun Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SY0298
Purification	ProA affinity purified
Applications	WB IHC ICC/IF
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Activator protein 1 antibody AP 1 antibody AP1 antibody cJun antibody Enhancer Binding Protein AP1 antibody Jun Activation Domain Binding Protein antibody JUN antibody Jun oncogene antibody JUN protein antibody Jun proto oncogene antibody JUN_HUMAN antibody JUNC antibody Oncogene JUN antibody p39 antibody Proto oncogene c jun antibody Proto oncogene cJun antibody Proto-oncogene c-jun antibody Transcription Factor AP 1 antibody Transcription factor AP-1 antibody Transcription Factor AP1 antibody V jun avian sarcoma virus 17 oncogene homolog antibody V jun sarcoma virus 17 oncogene homolog (avian) antibody V jun sarcoma virus 17 oncogene homolog antibody V-jun avian sarcoma virus 17 oncogene homolog antibody vJun Avian Sarcoma Virus 17 Oncogene Homolog antibody
Accession No.	Swiss-Prot#:P05412
Calculated MW	Predicted band size: 36 kDa
SDS-PAGE MW	Observed band size: 43 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:2000

IHC: 1:50-1:200

ICC/IF: 1:50-1:200

Images



All lanes: c-Jun Rabbit mAb at 1/1k dilution

Lane 1 : HUVEC whole cell lysates

Lane 2 : 293 whole cell lysates
Lane 3 : HepG2 whole cell lysates
Lane 4 : 3T3 whole cell lysates
Lane 5 : C6 whole cell lysates

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 36 kDa

Observed band size: 43 kDa

Exposure time: 5 seconds




All lanes : c-Jun Rabbit mAb at 1/1k dilution


Lane 1 : Wild-type HEK-293 cell lysate

Lane 2 : c-Jun knockdown HEK-293 cell lysate


Lysates/proteins at 20 µg per lane.




Formalin-fixed, paraffin-embedded human breast tissue stained for c-Jun using 48820 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded rat cerebral cortex tissue stained for c-Jun using 48820 at 1/100 dilution in immunohistochemical analysis.



Formalin-fixed, paraffin-embedded mouse skeletal muscle tissue stained for c-Jun using 48820 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence c-Jun antibody (48820) ICC/IF staining of c-Jun in NIH/3T3 cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 48820 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 488 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Background

Genes belonging to the Jun and Fos oncogene families encode nuclear proteins that are associated with a number of transcriptional complexes. The c-Jun protein is a major component of the transcription factor AP-1, originally shown to mediate phorbol ester tumor promoter (TPA)-induced expression of responsive genes through the TPA-response element (TRE). The Jun proteins form homo- and heterodimers which bind the TRE, while Fos proteins are active only as heterodimers with any of the Jun proteins. Fos/Jun heterodimers have a much higher affinity for the TRE than Jun homodimers. Ha-Ras augments c-Jun activity and stimulates phosphorylation of its activation domain. An inhibitor of Fos/Jun function, termed IP-1, associates with Fos and Jun and is inactivated upon phosphorylation induced by the cAMP-dependent protein kinase A (PKA).

Note: This product is for in vitro research use only and is not intended for use in humans or animals.