

CD31 Rabbit mAb

Catalog No: #48832

Package Size: #48832-1 50ul #48832-2 100ul



SAB[®]

Signalway Antibody

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	CD31 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	SU03-59
Purification	ProA affinity purified
Applications	WB, IHC, IP, FC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Adhesion molecule antibody CD31 antibody CD31 antigen antibody CD31 EndoCAM antibody EndoCAM antibody FLJ34100 antibody FLJ58394 antibody GPIIA antibody GPIIA' antibody PECA1 antibody PECA1_HUMAN antibody Pecam 1 antibody PECAM 1 CD31 EndoCAM antibody PECAM antibody PECAM-1 antibody Pecam1 antibody Platelet and endothelial cell adhesion molecule 1 antibody Platelet endothelial cell adhesion molecule antibody Platelet/endothelial cell adhesion molecule 1 antibody
Accession No.	Swiss-Prot#:P16284
Calculated MW	82kDa(Observed: 130 kDa)
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC:1:50-1:200 FC: 1:50-1:100


Images




Western blot analysis of CD31 on THP-1 cell lysates using anti-CD31 antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD31 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-CD31 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human uterus tissue using anti-CD31 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of THP-1 cells with CD31 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Cell adhesion molecules are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth and are thought to play an important role in embryogenesis and development. Neuronal cell adhesion molecule (NCAM) expression is observed in a variety of human tumors including neuroblastomas, rhabdomyosarcomas, Wilms tumors, Ewings sarcomas and some primitive myeloid malignancies. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the $\beta 2/\alpha 1$ and $\beta 2/\alpha M$ integrins. PECAM-1 (platelet/endothelial cell adhesion molecule-1), also referred to as CD31, is a glycoprotein expressed on the cell surfaces of monocytes, neutrophils, platelets and a subpopulation of T cells. VCAM-1 (vascular cell adhesion molecule-1) was first identified as an adhesion molecule induced on human endothelial cells by inflammatory cytokines such as IL-1, tumor necrosis factor (TNF) and lipopolysaccharide

(LPS). The KALIG gene encodes a nerve cell adhesion molecule (NCAM)-like protein and is deleted in 66% of patients with Kallmanns syndrome, anosmia with secondary hypogonadism.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.