

MEKK2 Rabbit mAb

Catalog No: #48987

Package Size: #48987-1 50ul #48987-2 100ul



**SAB**<sup>®</sup>

Signalway Antibody

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)

Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)


Description

|                       |                                                                                                                                                                                                                                                 |
|-----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name          | MEKK2 Rabbit mAb                                                                                                                                                                                                                                |
| Host Species          | Recombinant Rabbit                                                                                                                                                                                                                              |
| Clonality             | Monoclonal                                                                                                                                                                                                                                      |
| Clone No.             | SC68-02                                                                                                                                                                                                                                         |
| Purification          | ProA affinity purified                                                                                                                                                                                                                          |
| Applications          | WB, ICC, IHC, IP, FC                                                                                                                                                                                                                            |
| Species Reactivity    | Hu, Ms, Rt                                                                                                                                                                                                                                      |
| Immunogen Description | recombinant protein                                                                                                                                                                                                                             |
| Conjugates            | Unconjugated                                                                                                                                                                                                                                    |
| Other Names           | M3K2_HUMAN antibody Map3k2 antibody MAPK/ERK kinase kinase 2 antibody MEK kinase 2 antibody MEKK 2 antibody MEKK2b antibody Mitogen activated protein kinase kinase kinase 2 antibody Mitogen-activated protein kinase kinase kinase 2 antibody |
| Accession No.         | Swiss-Prot#:Q9Y2U5                                                                                                                                                                                                                              |
| Calculated MW         | 70 kDa                                                                                                                                                                                                                                          |
| Formulation           | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.                                                                                                                                                                            |
| Storage               | Store at -20°C                                                                                                                                                                                                                                  |


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of MEKK2 on different lysates using anti-MEKK2 antibody at 1/1,000 dilution. Positive control:  
Lane 1: HepG2      Lane 2: Rat brain Lane 3: SW480




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-MEKK2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-MEKK2 antibody. Counter stained with hematoxylin.




ICC staining MEKK2 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining MEKK2 in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining MEKK2 in SW480 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of HepG2 cells with MEKK2 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

## Background

Mitogen-activated protein (MAP) kinase cascades are activated by various extracellular stimuli including growth factors. The MEK kinases (also designated MAP kinase kinase kinases, MKKKs, MAP3Ks or MEKKs) phosphorylate and thereby activate the MEKs (also called MAP kinase kinases or MKKs), including ERK, JNK and p38. These activated MEKs in turn phosphorylate and activate the MAP kinases. The MEK kinases include Raf-1, Raf-B, Mos, MEK kinase-1, MEK kinase-2, MEK kinase-3, MEK kinase-4, ASK 1 (MEK kinase-5) and MAP3K6 (MEK kinase-6). MEK kinase-1 has been shown to phosphorylate MEK-1 via a Raf-independent pathway. Evidence suggests that MEK-3 is preferentially activated by MEK kinase-3 and that MEK-4 is activated by both MEK kinase-2 and MEK kinase-3. MEK kinase-4 has been shown to specifically activate the JNK pathway. ASK 1 activates both MEK-4 and MEK-3/MEK-6 pathways.

## References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.