## SIRT5 Rabbit mAb

Catalog No: #49271

Package Size: #49271-1 50ul #49271-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


# Description

Product Name	SIRT5 Rabbit mAb			
Host Species	Recombinant Rabbit			
Clonality	Monoclonal			
Clone No.	JJ084-01			
Purification	ProA affinity purified			
Applications	WB			
Species Reactivity	Hu, Ms			
Immunogen Description	recombinant protein			
Conjugates	Unconjugated			
Other Names	NAD dependent deacetylase sirtuin 5 antibody NAD dependent lysine demalonylase and desuccinylase sirtuin			
	5 mitochondrial antibody NAD dependent protein deacylase sirtuin 5 mitochondrial antibody NAD-dependent			
	protein deacylase sirtuin-5, mitochondrial antibody Regulatory protein SIR2 homolog 5 antibody Silent mating			
	type information regulation 2 S.cerevisiae homolog 5 antibody Sir2 like 5 antibody SIR2-like protein 5 antibody			
	SIR2L5 antibody SIR5_HUMAN antibody Sirt5 antibody Sirtuin 5 antibody Sirtuin type 5 antibody			
Accession No.	Swiss-Prot#:Q9NXA8			
Calculated MW	37/22 kDa			
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.			
Storage	Store at -20°C			

## **Application Details**

WB: 1:1,000-1:2,000

## **Images**



Western blot analysis of SIRT5 on Jurkat cells lysates using anti-SIRT5 antibody at 1/1,000 dilution.

# Background

SIRT5 is a human member of a family of proteins called Sirtuins (Sir2-like proteins) and are present in prokaryotes and eukaryotes. All Sir2-like

proteins have a sirtuin core domain, which contains a series of sequence motifs conserved in organisms ranging from bacteria to humans. Bacterial, yeast and mammalian sirtuins are able to metabolize NAD and possibly at as mono-ADP-ribosyltransferases. The enzymatic function of sirtuins is not yet completely understood but recent reports of histone-activated Sir2-mediated NAD metabolism and NAD-activated Sir2-mediated histone deacetylation suggest a possible coupled reciprocal activation mechanism involving interactions of Sir2 with NAD and the N epsilon-acetyl-lysine groups of acetylated histones.

$\overline{}$	- 1	 r۵		

Note: This product is for in vitro research use only and is not intended for use in humans or animals.