#### **Product Datasheet**

# HE4 Rabbit mAb

Catalog No: #49375

Package Size: #49375-1 50ul #49375-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


		4.5	
Des	cri	ntic	ı'n
レしこ	ווטפ	$\nu$ u $\epsilon$	/I I

Product Name	HE4 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JF62-09
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu, Ms
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	dJ461P17.6 antibody EDDM4 antibody epididymal protein 4 antibody Epididymal secretory protein E4
	antibody Epididymis specific whey acidic protein type four disulfide core antibody HE 4 antibody Major
	epididymis specific protein E4 antibody Major epididymis-specific protein E4 antibody MGC57529 antibody
	Putative protease inhibitor WAP5 antibody WAP 5 antibody WAP domain containing protein HE4 antibody
	WAP domain containing protein HE4 V4 antibody WAP four disulfide core domain 2 antibody WAP four
	disulfide core domain protein 2 antibody WAP four-disulfide core domain protein 2 antibody WAP5 antibody
	WFDC 2 antibody WFDC2 antibody WFDC2_HUMAN antibody
Accession No.	Swiss-Prot#:Q14508
Calculated MW	23 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


# **Application Details**

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


# **Images**




Western blot analysis of HE4 on different lysates using anti-HE4 antibody at 1/1,000 dilution. Positive control: Lane 1: SW480 Lane 2: SKOV-3




Immunohistochemical analysis of paraffin-embedded mouse lung tissue using anti-HE4 antibody. Counter stained with hematoxylin.



ICC staining HE4 in SKOV-3 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining HE4 in SW480 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of SW480 cells with HE4 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

# Background

HE4 (whey acidic protein (WAP)-type four-disulfide core-2, WFDC2) is a small secretory protein that may influence sperm maturation. HE4 gene expression is high in pulmonary epithelial cells and in some ovarian cancers. HE4 protein has a WAP motif that contains eight cysteines forming four disulfide bonds at the core of the protein. The WAP motif functions as a protease inhibitor in many of the family members that contain them.

### References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.				
The product of the first the recourse and only and to not interface for account furnished animals.				