## beta Tubulin(HRP conjugated) Rabbit mAb

Catalog No: #49382

Package Size: #49382-1 50ul #49382-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


		4.5	
Des	cri	ntic	ı'n
レしこ	011	$\nu$ u $\epsilon$	/I I

Product Name	beta Tubulin(HRP conjugated) Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JF41-50
Purification	ProA affinity purified
Applications	WB
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	HRP
Other Names	Beta 4 tubulin antibody Beta 5 tubulin antibody BetaTubulin antibody TBB5_HUMAN antibody TUBB
	antibody TUBB2 antibody TUBB2A antibody TUBB5 antibody tubulin beta 2A antibody Tubulin beta chain
	antibody Tubulin beta-5 chain antibody
Accession No.	Swiss-Prot#:P07437
Calculated MW	50 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

## **Application Details**

WB: 1:1,000-1:2,000

## **Images**



Western blot analysis of beta Tubulin on different lysates using anti-beta Tubulin antibody at 1/1,000 dilution. Positive control:

Lane 1: Hela Lane 2: PC-12 Lane 3: NIH/3T3

## Background

Tubulin is a major cytoskeleton component that has five distinct forms, designated  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  and e Tubulin.  $\alpha$  and  $\beta$  Tubulins form heterodimers which multimerize to form a microtubule filament. Multiple  $\beta$  Tubulin isoforms ( $\beta$ 1,  $\beta$ 2,  $\beta$ 3,  $\beta$ 4,  $\beta$ 5,  $\beta$ 6 and  $\beta$ 8) have been characterized and are expressed in mammalian tissues.  $\beta$ 1 and  $\beta$ 4 are present throughout the cytosol,  $\beta$ 2 is present in the nuclei and nucleoplasm, and  $\beta$ 3 is a neuron-specific

cytoskeletal protein.  $\gamma$  Tubulin forms the gammasome, which is required for nucleating microtubule filaments at the centrosome. Both  $\delta$  Tubulin and e Tubulin are associated with the centrosome.  $\delta$  Tubulin is a homolog of the Chlamydomonas  $\delta$  Tubulin Uni3 and is found in association with the centrioles, whereas e Tubulin localizes to the pericentriolar material. e Tubulin exhibits a cell-cycle-specific pattern of localization, first associating with only the older of the centrosomes in a newly duplicated pair and later associating with both centrosomes.

_						
Ю	$\sim$ t	$\sim$	-	n	റല	0
_	-	-		ш	-	•

Note: This product is for in vitro research use only and is not intended for use in humans or animals.