# FGF2 Rabbit mAb

Catalog No: #49431

Package Size: #49431-1 50ul #49431-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


# Description

Product Name	FGF2 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JM28-10
Purification	ProA affinity purified
Applications	WB, IP, FC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Basic fibroblast growth factor antibody Basic fibroblast growth factor bFGF antibody BFGF antibody FGF 2
	antibody FGF B antibody FGF-2 antibody Fgf2 antibody FGF2 basic antibody FGF2_HUMAN antibody FGFB
	antibody Fibroblast growth factor 2 (basic) antibody Fibroblast growth factor 2 antibody Fibroblast growth
	factor, basic antibody HBGF 2 antibody HBGF-2 antibody HBGF2 antibody HBGH 2 antibody HBGH2
	antibody Heparin binding growth factor 2 precursor antibody Heparin-binding growth factor 2 antibody
	Prostatropin antibody
Accession No.	Swiss-Prot#:P09038
Calculated MW	31 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

# **Application Details**

WB: 1:500-1:1000IP: 1:50FC: 1:50-1:100

### **Images**



Flow cytometric analysis of Jurkat cells with FGF2 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

# Background

Fibroblast growth factor-1 (FGF-1), also designated acidic FGF, and fibroblast growth factor-2 (FGF-2), also designated basic FGF, are members of a family of growth factors that stimulate proliferation of cells of mesenchymal, epithe-lial and neuroectodermal origin. Additional members of the FGF family include the oncogenes FGF-3 (Int2) and FGF-4 (hst/Kaposi), FGF-5, FGF-6, FGF-7 (KGF), FGF-8 (AIGF), FGF-9 (GAF) and FGF-10-FGF-23. Members of the FGF family share 30-55% amino acid sequence identity and similar gene structure, and are capable of transforming cultured cells when overexpressed in transfected cells. Cellular receptors for FGFs are members of a second multigene family including four tyrosine kinases, designated Flg (FGFR-1), Bek (FGFR-L), TKF and FGFR-3.

#### References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.