

Lipocalin-2 Rabbit mAb

Catalog No: #49452

Package Size: #49452-1 50ul #49452-2 100ul



Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	Lipocalin-2 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JM10-67
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	24p3 antibody 25 kDa alpha 2 microglobulin related subunit of MMP9 antibody 25 kDa alpha-2-microglobulin-related subunit of MMP-9 antibody Alpha 2 microglobulin related protein antibody HGNC:6526 antibody HNL antibody Lcn 2 antibody Lcn2 antibody Lipocalin-2 antibody Migration stimulating factor inhibitor antibody MSFI antibody Neutrophil gelatinase associated lipocalin antibody Neutrophil gelatinase associated lipocalin precursor antibody Neutrophil gelatinase-associated lipocalin antibody NGAL antibody NGAL_HUMAN antibody Oncogene 24p3 antibody Oncogenic lipocalin 24p3 antibody p25 antibody Siderocalin antibody siderocalin LCN2 antibody SV40 induced 24P3 protein antibody Uterocalin antibody
Accession No.	Swiss-Prot#:P80188
Calculated MW	23 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-5,000IHC: 1:50-1:200 ICC: 1:100-1:500FC: 1:50-1:100


Images




Western blot analysis of Lipocalin-2 on A431 cells lysates using anti-Lipocalin-2 antibody at 1/500 dilution.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue using anti-Lipocalin-2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-Lipocalin-2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human lung tissue using anti-Lipocalin-2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human uterus tissue using anti-Lipocalin-2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human pancreas tissue using anti-Lipocalin-2 antibody. Counter stained with hematoxylin.



ICC staining Lipocalin-2 in 293T cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Lipocalin-2 in Hela cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Lipocalin-2 in SW480 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

In addition to the monomeric mammalian progelatinase, two additional forms of progelatinase have been identified. The shorter of these additional forms is a covalently linked, disulfide-bridged protein that heterodimerizes with a short protein; an α 2-Microglobulin-related protein known as neutrophil gelatinase-associated lipocalin (NGAL), which is moderately expressed in breast and lung tissues. NGAL belongs to the lipocalin family and has a high degree of similarity with rat α 2-Microglobulin-related protein and mouse protein 24p3. NGAL is able to bind a derivative of the bacterial chemotactic peptide, suggesting that it has important immuno-modulatory functions. NGAL has been described as an inflammatory protein; it is released into the circulation as a result of the inflammatory activation of leukocytes initiated by the extra-corporeal circulation. In addition, NGAL synthesis is induced in epithelial cells in inflammatory and neoplastic colorectal diseases. In conclusion, NGAL may serve as a scavenger of bacterial products to function in the anti-inflammatory process.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.