Ephrin B2 Rabbit mAb

Catalog No: #49626

Package Size: #49626-1 50ul #49626-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


			4.5	
	Δcc	rır	\TIC	۱n
u	esc	ш	ıιι	ווע

Storage	Store at +4°C for short term. Store at -20°C for long term. Avoid freeze/thaw cycle.		
Formulation	Rabbit IgG in 10mM phosphate buffered saline, pH 7.4, 150mM sodium chloride, 0.05% BSA, 0.02% sodium azide and 50% glycerol.		
Calculated MW	50 kDa		
Accession No.	Swiss-Prot#:P52799		
	OTTMUSP00000024973 antibody		
	eph related kinase 5 antibody MGC126226 antibody MGC126227 antibody MGC126228 antibody		
	antibody HTK-L antibody HTKL antibody LERK 5 antibody LERK-5 antibody LERK5 antibody Ligand of		
	Ephrin-B2 antibody EphrinB2 antibody EPLG 5 antibody EPLG5 antibody Htk L antibody HTK ligand		
	kinase ligand 5 antibody EPH-related receptor tyrosine kinase ligand 5 antibody ephrin B2 antibody		
Other Names	EFN B2 antibody EFNB 2 antibody Efnb2 antibody EFNB2_HUMAN antibody Eph related receptor tyrosine		
Conjugates	Unconjugated		
Immunogen Description	A synthesized peptide derived from human Ephrin B2		
Species Reactivity	Hu, Ms, Rt		
Applications	WB, ICC/IF		
Purification	Affinity-chromatography		
Clone No.	JM53-21		
Clonality	Monoclonal		
Product Name	Ephrin B2 Rabbit mAb		


Application Details

WB 1:1000-1:2000 ICC/IF 1:50-1:200


Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of Ephrin B2 expression in (1) HEK293 cell lysate; (2) Mouse spleen lysate.



Immunofluorescent analysis using the Antibody at 1:50 dilution.

Background

Cell surface transmembrane ligand for Eph receptors, a family of receptor tyrosine kinases which are crucial for migration, repulsion and adhesion during neuronal, vascular and epithelial development. Binds promiscuously Eph receptors residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.