

USP11 Rabbit mAb

Catalog No: #49628

Package Size: #49628-1 50ul #49628-2 100ul



SAB[®]
Signalway Antibody

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	USP11 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal
Clone No.	JM33-62
Purification	ProA affinity purified
Applications	WB, ICC/IF, IP, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Conjugates	Unconjugated
Other Names	Deubiquitinating enzyme 11 antibody Ubiquitin carboxyl-terminal hydrolase 11 antibody Ubiquitin carboxyl-terminal hydrolase X linked antibody ubiquitin specific peptidase 11 antibody Ubiquitin specific protease 11 antibody Ubiquitin thiolesterase 11 antibody Ubiquitin-specific-processing protease 11 antibody UBP11_HUMAN antibody UHX1 antibody USP11 antibody
Accession No.	Swiss-Prot#:P51784
Calculated MW	110 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:1,000ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of USP11 on 293T cell using anti-USP11 antibody at 1/1,000 dilution.




ICC staining USP11 in A549 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining USP11 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining USP11 in SH-SY5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Jurkat cells with USP11 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black).

Background

The ubiquitin (Ub) pathway involves three sequential enzymatic steps that facilitate the conjugation of Ub and Ub-like molecules to specific protein substrates. Through the use of a wide range of enzymes that can add or remove ubiquitin, the Ub pathway controls many intracellular processes such as signal transduction, transcriptional activation and cell cycle progression. USP11 (ubiquitin specific peptidase 11), also known as UHX1, is a 920 amino acid deubiquitinating enzyme that participates in the Ub pathway. Localized to the nucleus, USP11 associates with both Ran BP-M (Ran binding protein M) and with the tumor suppressor BRCA2. Through these associations, USP11 functions to either inhibit ubiquitination of these proteins or to remove ubiquitin residues that have already been attached to these proteins. USP11 is implicated in several X-linked retinal diseases and, due to its ability to deubiquitinate BRCA2, may play a role in tumor suppression.

References

Note: This product is for in vitro research use only and is not intended for use in humans or animals.