

Cyclin A2 Rabbit Polyclonal Antibody

Catalog No: #55266

Package Size: #55266-1 50ul #55266-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Cyclin A2 Rabbit Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Isotype	IgG
Purification	Affinity purification
Applications	WB,IHC,IF
Species Reactivity	Human,Mouse,Rat
Immunogen Description	A synthetic peptide of human Cyclin A2
Conjugates	Unconjugated
Other Names	CCN1;CCNA;Cyclin A2;CCNA2;cyclin-A2
Accession No.	Swiss Prot:P20248GenelD:890
Calculated MW	48kDa
SDS-PAGE MW	55kDa
Formulation	Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
Storage	Store at -20°C. Avoid freeze / thaw cycles.


Application Details

WB□1:500 - 1:2000IHC□1:50 - 1:200IF□1:50 - 1:200


Images




Immunohistochemistry of paraffin-embedded human well-differentiated squamous skin carcinoma using Cyclin A2 at dilution of 1:100 (40x lens).



Immunohistochemistry of paraffin-embedded mouse kidney using Cyclin A2 at dilution of 1:100 (40x lens).



Immunofluorescence analysis of U2OS cells using Cyclin A2 at dilution of 1:100. Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using Cyclin A2 at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit (RM00021). Exposure time: 90s.

Background

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. In contrast to cyclin A1, which is present only in germ cells, this cyclin is expressed in all tissues tested. This cyclin binds and activates CDC2 or CDK2 kinases, and thus promotes both cell cycle G1/S and G2/M transitions.

Published Papers

el at., CD155 Cooperates with PD-1/PD-L1 to Promote Proliferation of Esophageal Squamous Cancer Cells via PI3K/Akt and MAPK Signaling Pathways. In *Cancers (Basel)* on 2022 Nov 15 by Xiyang Tan, Jie Yang, et al.. PMID:36428703, (2022)

[PMID:36428703](https://pubmed.ncbi.nlm.nih.gov/36428703/)

Note: This product is for in vitro research use only and is not intended for use in humans or animals.