## CDKN2C Rabbit mAb

Catalog No: #56630

Package Size: #56630-1 50ul #56630-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


## Description

| Product Name          | CDKN2C Rabbit mAb  |
|-----------------------|--|
| Host Species          | Rabbit   |
| Clonality             | Monoclonal   |
| Isotype               | Rabbit IgG   |
| Purification          | Affinity-chromatography  |
| Applications          | WB IHC ICC/IF IP   |
| Species Reactivity    | Human Mouse Rat  |
| Specificity           | CDKN2C Antibody detects endogenous levels of total CDKN2C  |
| Immunogen Description | A synthesized peptide derived from human CDKN2C  |
| Conjugates            | Unconjugated   |
| Other Names           | Cdkn2c; CDKN6; INK4C; p18; p18 inhibits CDK4; p18 INK4c; p18 INK6; p18-INK4c; p18-INK6;            |
| Accession No.         | Uniprot:P42773   |
| Calculated MW         | 15kDa  |
| Formulation           | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage               | Store at +4Λ C short term. Store at -20Λ C long term. Avoid freeze / thaw cycle.                   |

## **Application Details**

WB:1:500~1:2000
IHC:1:50~1:200
ICC/IF:1:50~1:200
IP:1:40

## **Images**



All lanes: CDKN2C Rabbit mAb at 1/1k dilution

Lane 1 : MCF7 whole cell lysates Lane 2 : SH-SY5Y whole cell lysates Lane 3 : Mouse bone lysates Lane 4 : Rat bone lysates

ysates


Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) at 1/20000 dilution

Predicted band size: 18 kDa Observed band size: 16 kDa

Exposure time: 8 seconds




All lanes:CDKN2C Rabbit mAb at 1/1k dilution


Lane 1: Wild-type Hela cell lysate

Lane 2: CDKN2C knockdown Hela cell lysate

Lysates/proteins at 20 µg per lane.



Formalin-fixed, paraffin-embedded human breast cancer tissue stained for CDKN2C using 56630 at 1/100 dilution in immunohistochemical analysis.



Immunocytochemistry/ Immunofluorescence CDKN2C antibody (56630) ICC/IF staining of CDKN2C in Hela cells. Cells were fixed with 4% Paraformaldehyde permeabilized with 0.1% Triton X-100.

Samples were incubated with 56630 at a working dilution of 1/100. The secondary antibody was Alexa FluorB 647 goat anti rabbit, used at a dilution of 1/500.

Nuclei were counterstained with DAPI.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.