Ube2N / Ubc13 Rabbit mAb

Catalog No: #59831

Package Size: #59831-1 50ul #59831-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


Description

| Product Name | Ube2N / Ubc13 Rabbit mAb |
|-----------------------|--|
| Host Species | Rabbit |
| Clonality | Monoclonal |
| Isotype | Rabbit IgG |
| Purification | Affinity-chromatography |
| Applications | WB IHC ICC/IF |
| Species Reactivity | Human Mouse Rat |
| Specificity | Ube2N / Ubc13 Antibody detects endogenous levels of total Ube2N / Ubc13 |
| Immunogen Description | A synthesized peptide derived from human Ube2N / Ubc13 |
| Other Names | BLU; HEL-S-71; Ubc13; UbCH ben; UbcH13; UBCHBEN; Ube2n; |
| Accession No. | Uniprot:P61088 |
| Formulation | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

Application Details

WB 1:500~1:2000 IHC 1:50~1:200 ICC/IF 1:50~1:200

Images



Western blot analysis of Ube2N / Ubc13 expression in Daudi cell lysate.

Background

The UBE2V1-UBE2N and UBE2V2-UBE2N heterodimers catalyze the synthesis of non-canonical 'Lys-63'-linked polyubiquitin chains. This type of polyubiquitination does not lead to protein degradation by the proteasome. Mediates transcriptional activation of target genes. Plays a role in the control of progress through the cell cycle and differentiation.

| Note: This product is for in vitro research use only and is not intended for use in humans or animals. | | | |
|--|--|--|--|
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |