Product Datasheet

Hepatitis C Virus NS3 antibody

69kDa

1mg/ml

Catalog No: #62137

Package Size: #62137 100ul



Support: tech@signalwayantibody.com

Description Hepatitis C Virus NS3 antibody **Product Name Host Species** Rabbit Clonality Polyclonal Purification The antibody was affinity purified by Protein A. **ELISA** Applications Species Reactivity Hepatitis C Virus Immunogen Type Peptide Immunogen Description peptide derived from HCV-NS3: 1451-1550/3010 **Target Name** Hepatitis C Virus NS3

Liquid in 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Store at +4°C for short term. Store at -20°C for long term. Avoid freeze/thaw cycle.

Application Details

ELISA 1:5000-10000

Calculated MW

Concentration

Formulation

Storage

Background

Core protein packages viral RNA to form a viral nucleocapsid, and promotes virion budding. Modulates viral translation initiation by interacting with HCV IRES and 40S ribosomal subunit. Also regulates many host cellular functions such as signaling pathways and apoptosis. Prevents the establishment of cellular antiviral state by blocking the interferon-alpha/beta (IFN-alpha/beta) and IFN-gamma signaling pathways and by inducing human STAT1 degradation. Plays an important role in virus-mediated cell transformation leading to hepatocellular carcinomas. Interacts with, and activates STAT3 leading to cellular transformation. May repress the promoter of p53, and sequester CREB3 and SP110 isoform3/Sp110b in the cytoplasm. Also represses cell cycle negative regulating factor CDKN1A, thereby interrupting an important check point of normal cell cycle regulation. Targets transcription factors involved in the regulation of inflammatory responses and in the immune response: suppresses NK-kappaB activation, and activates AP-1. Mediates apoptotic pathways throught association with TNF-type receptors TNFRSF1A and LTBR, although its effect on death receptors-induced apoptosis remains controvertial. Enhances TRAIL mediated apoptosis, suggesting that it might play a role in immune-mediated liver cell injury. Secreted core protein is able to bind C1QR1 at the T-cell surface, resulting in down-regulation of T-lymphocytes proliferation. May transactivate human MYC, Rous sarcoma virus LTR, and SV40 promoters. May suppress the human FOS and HIV-1 LTR activity. May alter lipid metabolism by interacting with hepatocellular proteins involved in lipid accumulation and storage.

Note: This product is for in vitro research use only and is not intended for use in humans or animals.